



Ideals of Regular Operators on l_2

Author(s): W. Arendt and A. R. Sourour

Source: *Proceedings of the American Mathematical Society*, Vol. 88, No. 1 (May, 1983), pp. 93-96

Published by: American Mathematical Society

Stable URL: <http://www.jstor.org/stable/2045116>

Accessed: 04/03/2009 15:05

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/page/info/about/policies/terms.jsp>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/action/showPublisher?publisherCode=ams>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is a not-for-profit organization founded in 1995 to build trusted digital archives for scholarship. We work with the scholarly community to preserve their work and the materials they rely upon, and to build a common research platform that promotes the discovery and use of these resources. For more information about JSTOR, please contact support@jstor.org.



American Mathematical Society is collaborating with JSTOR to digitize, preserve and extend access to *Proceedings of the American Mathematical Society*.

<http://www.jstor.org>

IDEALS OF REGULAR OPERATORS ON l^2

W. ARENDT AND A. R. SOUROUR

ABSTRACT. Let \mathcal{L}^r be the Banach algebra (and Banach lattice) of all regular operators on l^2 , i.e. the algebra of all operators A on l^2 which are given by a matrix (a_{mn}) such that $(|a_{mn}|)$ defines a bounded operator $|A|$. We show that there exists exactly one nontrivial closed subspace of \mathcal{L}^r which is both a lattice-ideal and an algebra-ideal of \mathcal{L}^r , namely the space $\mathcal{K}^r = \{A \in \mathcal{L}^r: |A| \text{ is compact}\}$. We also show that every nontrivial ideal in \mathcal{L}^r is included in \mathcal{K}^r .

It is well known that the only nontrivial closed (algebra) ideal in the Banach algebra $\mathcal{L}(H)$ of all bounded operators on a separable Hilbert space H is the ideal $\mathcal{K}(H)$ consisting of all compact operators on H and that $\mathcal{K}(H)$ includes every nontrivial ideal. In this note we prove order-theoretic analogues of these results.

To give a precise statement we need some notation. Let $H = l^2$, the Hilbert space of all (real or complex) square-summable sequences, and denote by $\{e_n\}$, $n = 1, 2, \dots$, the standard basis in l^2 . Every bounded operator A on l^2 can be represented by a matrix (a_{mn}) given by $a_{mn} = (Ae_n | e_m)$, $m, n = 1, 2, \dots$, where $(f | g)$ denotes the scalar product of $f, g \in l^2$. If the matrix $(|a_{mn}|)$ also defines a bounded operator $|A|$, then A is called *regular*, and $|A|$ is called the *modulus* of A . For example, every operator of finite rank is obviously regular, but there exist compact operators which are not regular [4, IV, §1, Example 2]. We denote by \mathcal{L}^r the space of all regular operators on l^2 . It is known that \mathcal{L}^r is a subalgebra of $\mathcal{L}(l^2)$ and is a Banach algebra under the r -norm $\|A\|_r$, defined as the operator norm of $|A|$, i.e.

$$\|A\|_r = \||A|\|$$

where $\|T\|$ denotes the operator norm of T (cf. [4, IV, §1]). A bounded operator A on l^2 is called *positive* (we write $A \geq 0$) if $(Ae_n | e_m) \geq 0$ for all natural numbers m, n . With this order relation, \mathcal{L}^r becomes a Banach lattice (cf. [4, IV, §1]). A *lattice-ideal* of \mathcal{L}^r is, by definition, a subspace \mathcal{J} of \mathcal{L}^r such that if $A \in \mathcal{J}$, $B \in \mathcal{L}^r$, $|B| \leq |A|$, then $B \in \mathcal{J}$. For the purpose of this paper, we say briefly that \mathcal{J} is an *ideal* if it is both an algebra-ideal and a lattice-ideal.

A regular operator is called *r-compact* if A can be approximated in the r -norm by operators of finite rank. The space \mathcal{K}^r of all r -compact operators is a closed ideal in \mathcal{L}^r . Our main theorem gives a confirmation of the following conjecture by H. H. Schaefer: \mathcal{K}^r is the only closed nontrivial ideal in \mathcal{L}^r . Furthermore, we show that every nontrivial ideal is included in \mathcal{K}^r .

Received by the editors June 7, 1982.

1980 *Mathematics Subject Classification*. Primary 47D30; Secondary 47B55, 47B05.

©1983 American Mathematical Society
0002-9939/82/0000-1119/\$01.75

For vectors $f, g \in l^2$, we denote by $f \otimes g$ the rank one operator given by

$$(f \otimes g)h = (h|f)g \quad (h \in l^2).$$

The orthogonal projection onto $\text{span}\{e_1, \dots, e_n\}$ will be denoted by P_n (i.e., $P_n = \sum_{m=1}^n e_m \otimes e_m$) and will be called an *initial projection*. Note that $P_n \geq 0$, $(1 - P_n) \geq 0$ for every $n \in N$ (the set of natural numbers).

The following proposition is well known and easy to prove.

PROPOSITION 1. *Let $A \in \mathcal{L}(l^2)$. Then*

- (i) $\|A\| = \sup_{n \in N} \|P_n A P_n\| = \lim_{n \rightarrow \infty} \|P_n A P_n\|$;
- (ii) A is compact if and only if $\lim_{n \rightarrow \infty} \|(1 - P_n)A(1 - P_n)\| = 0$.

We give a characterization of r -compact operators (cf. [4, p. 293, Corollary 1]).

PROPOSITION 2. (i) *Let $B, C \in \mathcal{L}^r$, $|B| \leq C$. If C is compact, then $B \in \mathcal{K}^r$.*

(ii) $\mathcal{K}^r = \{A \in \mathcal{L}^r: |A| \text{ is compact}\}$.

(iii) \mathcal{K}^r is a closed ideal in \mathcal{L}^r .

PROOF. Let $F_n = P_n B P_n + (1 - P_n) B P_n + P_n B (1 - P_n)$. Then F_n is an operator of finite rank and

$$|B - F_n| = |(1 - P_n)B(1 - P_n)| \leq (1 - P_n)|B|(1 - P_n) \leq (1 - P_n)C(1 - P_n).$$

It follows from Proposition 1 (ii) that $\lim_{n \rightarrow \infty} \|B - F_n\|_r = 0$. Hence $B \in \mathcal{K}^r$. This proves part (i).

From (i), it follows that \mathcal{K}^r includes $\{A \in \mathcal{L}^r: |A| \text{ compact}\}$. To prove the reverse inclusion, first observe that $|F|$ is compact if F is an operator of finite rank (this follows from (i) since if $F = \sum_{i=1}^n x_i \otimes y_i$, then $|F| \leq \sum_{i=1}^n |x_i| \otimes |y_i|$). Now, let $A \in \mathcal{K}^r$. Then $\lim_{n \rightarrow \infty} \|A - F_n\|_r = 0$ for a sequence $\{F_n\}$ of finite rank operators. This implies that $\lim_{n \rightarrow \infty} \||A| - |F_n|\| = 0$. Therefore $|A|$ is a limit of a sequence of compact operators and so is compact.

Part (iii) follows immediately from (i) and (ii). \square

THEOREM 1. *Let \mathcal{G} be a proper ideal in \mathcal{L}^r . Then $\mathcal{G} \subset \mathcal{K}^r$.*

PROOF. Let \mathcal{G} be an ideal in \mathcal{L}^r which is not included in \mathcal{K}^r . We will show that $\mathcal{G} = \mathcal{L}^r$.

There exists a positive operator A in \mathcal{G} which is not compact. Let P_{n_1} be an initial projection such that $\|P_{n_1} A P_{n_1}\| \geq \frac{1}{2} \|A\|$. By induction, we can find a sequence of increasing initial projections $\{P_{n_j}\}$, $j = 1, 2, \dots$, such that

$$\|(P_{n_{j+1}} - P_{n_j})A(P_{n_{j+1}} - P_{n_j})\| \geq \frac{1}{2} \|(1 - P_{n_j})A(1 - P_{n_j})\|.$$

Since A is not compact, the norms $\|(1 - P_{n_j})A(1 - P_{n_j})\|$ are bounded below by a positive number δ . Let $B_j = (P_{n_{j+1}} - P_{n_j})A(P_{n_{j+1}} - P_{n_j})$, so $\|B_j\| \geq \delta/2$. The operator $B = \sup_{j \in N} B_j$ belongs to \mathcal{G} since $0 \leq B \leq A$. Denote by H_j the range of $P_{n_{j+1}} - P_{n_j}$ and let g_j be a positive vector in H_j of norm 1 such that $\|B_j g_j\| \geq \delta/2$. Let $R_n = e_1^n \otimes g_n$, where e_1^n is the first basis vector in H_n (i.e., $e_1^k = e_{n_k+1}$), and let $R = \sup_{n \in N} R_n$. Thus R is a positive operator of norm 1. Let $M_n = B_n R_n = e_1^n \otimes B_n g_n$ and let $M = \sup_{n \in N} M_n$, so $M = BR \in \mathcal{G}$. Finally, let $T = M^* M$. Then $T \in \mathcal{G}$ and

$T = \sup_{n \in N} T_n$, where $T_n = M_n^* M_n$. So T_n is represented (with respect to the standard basis) by the diagonal matrix $\sum \alpha_n e_1^n \otimes e_1^n$, where $\alpha_n = \|B_n g_n\|^2$. There is a permutation matrix U such that

$$UTU = \text{diag}(\alpha_1, 0, \alpha_2, 0, \alpha_3, \dots).$$

The sequence $\{\alpha_n\}$ is bounded below, therefore the matrix

$$\text{diag}(\alpha_1^{-1}, 0, \alpha_2^{-1}, 0, \alpha_3^{-1}, 0, \dots)$$

defines a bounded positive operator D , and so $F = DUTU = \text{diag}(1, 0, 1, 0, \dots)$ belongs to \mathcal{F} . If V is the permutation operator defined by $Ve_{2n-1} = e_{2n}$, $Ve_{2n} = e_{2n-1}$ ($n \in N$), then $1 = F + V F V \in \mathcal{F}$. Therefore $\mathcal{F} = \mathcal{L}^r$. \square

LEMMA. Every nonzero algebra ideal in \mathcal{L}^r contains the finite rank operators.

PROOF. (This is the same proof as the well-known proof of the analogous result in $\mathcal{L}(l^2)$; we merely point out that all the operators involved are regular.) Let \mathcal{F} be a nonzero algebra ideal in \mathcal{L}^r and let A be a nonzero operator in \mathcal{F} . Pick a vector f_1 such that $\|A f_1\| = 1$. For two arbitrary vectors f and g , we have $f \otimes g = (A f_1 \otimes g)A(f \otimes f_1)$, and so every operator of rank one belongs to \mathcal{F} . Finally, every finite rank operator is a sum of rank one operators and therefore belongs to \mathcal{F} . \square

As an immediate consequence of Theorem 1 and the Lemma we obtain the main result.

THEOREM 2. \mathcal{K}^r is the only nontrivial closed ideal in \mathcal{L}^r .

COROLLARY. $\mathcal{K}^r = \{A \in \mathcal{L}^r: \lim_{n \rightarrow \infty} |A| R e_n = 0 \text{ for every positive operator } R\}$.

REMARK 1. There are nontrivial closed lattice-ideals as well as nontrivial closed algebra-ideals in \mathcal{L}^r other than \mathcal{K}^r . For example, $\{T \in \mathcal{L}^r: \lim_{n, m \rightarrow \infty} (T e_n | e_m) = 0\}$ is a nontrivial closed lattice-ideal different from \mathcal{K}^r . Moreover, there exists a compact operator $A \in \mathcal{L}^r$ such that $|A|$ is not compact [4, IV, §1, Example 2]. Hence $\mathcal{K}(l^2) \cap \mathcal{L}^r = \{T \in \mathcal{L}^r: T \text{ is compact}\}$ is a nontrivial closed algebra-ideal in \mathcal{L}^r which contains \mathcal{K}^r properly.

REMARK 2. By Theorem 1, \mathcal{K}^r is the largest ideal in \mathcal{L}^r . The smallest ideal \mathcal{F}_s is the one which is generated by the finite rank operators (by the Lemma). Thus it has the form $\mathcal{F}_s = \{T \in \mathcal{L}^r: |T| \leq h \otimes h \text{ for some positive } h \in l^2\}$. This ideal \mathcal{F}_s is properly included in the ideal of Hilbert-Schmidt operators, which, in turn, is properly included in \mathcal{K}^r .

REMARK 3. The situation is different if we replace l^2 by $L^2[0, 1]$. For the definition of $\mathcal{L}^r(L^2)$, see [4, IV, §1]. The space $\mathcal{K}^r(L^2)$ of all regular operators on L^2 which can be approximated in the r -norm by operators of finite rank is a closed ideal in $\mathcal{L}^r(L^2)$ (this follows from [4, p. 293, Corollary 2]); and by the same proof that we used for the Lemma, it is the smallest closed ideal in $\mathcal{L}^r(L^2)$. The space $\mathcal{K}_1 = \{T \in \mathcal{L}^r(L^2); |T| \text{ is compact}\}$ is obviously closed, and it follows from the Dodds-Fremlin Theorem ([3], see also [1]) that \mathcal{K}_1 is a lattice-ideal. This implies that it is also an algebra-ideal. However, \mathcal{K}_1 includes $\mathcal{K}^r(L^2)$ properly (see [2, 3.7 in connection with 2.6]). Another nontrivial closed ideal in $\mathcal{L}^r(L^2)$ is \mathcal{F} , the space of all regular integral

operators (or kernel operators), i.e. operators T of the form

$$Tf(s) = \int k(t, s)f(t) dt,$$

where k is a measurable function such that $|k|$ is the kernel of a bounded operator. (The ideal \mathcal{G} is $(L^2 \otimes L^2)^{\perp\perp}$, the band in $\mathcal{L}'(L^2)$ generated by the finite rank operators [4, IV, 9.8].) In the atomic case ($H = l^2$), all regular operators are kernel operators. Here, in the nonatomic case, \mathcal{G} is a proper closed ideal of $\mathcal{L}'(L^2)$, properly includes $\mathcal{K}'(L^2)$, and is not comparable to \mathcal{K}_1 . Furthermore $\mathcal{K}'(L^2) = \mathcal{K}_1 \cap \mathcal{G}$ [4, p. 293, Corollary 2]. In view of this, one nonatomic version of the problem solved in Theorem 2 is the following question: Is $\mathcal{K}'(L^2)$ the only nontrivial closed ideal in the Banach lattice algebra of kernel operators on L^2 ?

REFERENCES

1. C. D. Aliprantis and O. Burkinshaw, *Positive compact operators on Banach lattices* Math. Z. **174** (1980), 289–298.
2. W. Arendt, *On the σ -spectrum of regular operators and the spectrum of measures*, Math. Z. **179** (1981), 271–287.
3. P. Dodds and D. H. Fremlin, *Compact operators in Banach lattices*, Israel J. Math. **34** (1979), 287–320.
4. H. H. Schaefer, *Banach lattices and positive operators*, Springer-Verlag, Berlin-Heidelberg-New York, 1974.

MATHEMATISCHES INSTITUT DER UNIVERSITÄT, 74 TÜBINGEN, FEDERAL REPUBLIC OF GERMANY

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF VICTORIA, VICTORIA, BRITISH COLUMBIA, CANADA V8W 2Y2