## **TUM-IAS Primary Sources** Essays in Technology and Science

Volume 1, Issue 1, November 2011

#### 

#### This Issue: Non-Gaussian Stochastic Models



Foreword 1



Featured Essay: CARMA processes driven by non-Gaussian noise: Robert J. Stelzer 5

Author profile <u>42</u>

About Primary Sources 46



# Getting a grip on processes that govern the world

Presenting Robert Stelzer's Primary Sources essay

# Stochastic processes govern the world!

Sometimes we are rudely awakened to this fact when a really catastrophic event happens such as an earthquake with unexpected strength at an unexpected location.

That is the gist of non-traditional stochastic modeling: accounting for "unexpected events." Although such events cannot be predicted, they can be modeled.

Stochastic modeling has been at the heart of science and engineering for a long time, starting with the work of Pascal on games and then Gauss on

measurement errors, leading to the binomial distribution and in the limit, the Gaussian distribution. Many random physical phenomena are indeed governed by such distributions, a lot of theory has been developed for them and they have invaded engineering ranging from the design of electronic amplifiers and telecommunication links to helicopter control via Kalman filtering. Somehow, and thanks to the famous (Lindeberg-Lévy) central limit theorem, there is a belief that when you have the sum of a large number of presumably small effects, you automatically get a near-Gaussian distribution and that therefore all stochastic disturbances in engineering can be meaningfully modeled by Gaussian models...



Technische Universität München

Presenting Robert Stelzer's Primary Sources Essay

### The truth, however, is much more complex. Many natural and human-made probabilistic phenomena do not adhere to a Gaussian distribution.

This is already pretty evident from the properties of the latter. It has a very pronounced average value; the distribution is symmetric around that value and tapers off exponentially. Three or four "sigmas" away (sigma is a number equal to the standard deviation of the random variable) and the probability that the process has such a value is virtually nil. Already early on in the development of probability theory, the famous French mathematician Lévy defined a continuous distribution function for a new class of positive random variables, characteristic of some physical effects in spectroscopy and deviating from the Gaussian case. That new distribution has then been found useful for the modeling of economic effects such as the return on shares in the stock market. Pareto drew the attention of the economic community to these so-called "power law probability distributions," a class of distributions that are described by a power law and hence have a "fat tail," leading to statements such as "80% of world's wealth is in 20% of the hands."

It seems that many meaningful distributions obey such laws: e.g., size

of islands, Internet traffic, 1/f noise, size of sand particles, you name it – reason enough to give these distributions a great deal of attention!

Probability theorists generalized the distributions discovered by Lévy and Pareto and defined a class that they called "stable distributions." By this they meant that a sum of two random variables with such a distribution has again such a distribution.

This then appears to be an eminently useful class for modeling a great many basic phenomena in nature, engineering and economics, greatly generalizing the work done with Gaussian distributions (stable as well, but in a very special case). Very early on, however, problems with this approach appeared. It turns out that there is no closed-form expression for a general stable distribution, even though its so-called "characteristic function" (the Fourier transform of its distribution) is fully defined by four numerical quantities. Equally serious is the problem of determining the resulting distribution when a random variable with a stable distribution is driving a process ("is filtered," one says).

Foreword

**Featured Essay** 

**Author Profile** 

**Primary Sources** 

About

**Featured Essay** 

Foreword

**Author Profile** 

About **Primary Sources** 

The need to address the latter, fundamental problem crystallized one of the first Focus Groups at the TUM Institute for Advanced Study. Under the inspired leadership of Prof. Claudia Klüppelberg the problem of modeling systems driven by stable distributions was tackled both fundamentally and in relation to a variety of potential applications. Dr. (now Prof.) Robert Stelzer studied the filtering problem in depth, and the results of his investigations are described in this issue of Primary Sources. The results should be useful for a wide range of interesting scientific and engineering cases. Distributions do not come alone; they are integral parts of an overall system. A wind

turbine, for instance, is a dynamical system that is being driven by wind stochastics. Nanoscale circuits undergo fluctuations that influence

The point of view taken by Robert Stelzer is characteristic for a signal processing approach. He models the underlying system by a so-called CARMA filter - the horrible acronym standing for Continuous-time Auto Regressive Moving Average system.

This is a pretty general approach, applicable to linear time-invariant systems evolving in continuous time. From basic analysis we know that functions on an interval can often be represented by a ratio of polynomials (e.g., a rational approximation). When one does that to a transfer function in continuous time, one obtains a CARMA filter. The situation is then viewed as a filter processing a random

> process and producing a stochastic output. In many situations encountered in practice, one wishes to identify both the filter coefficients and the

their output. In such cases the basic system that is being driven can be modeled from its physics. Less clear is the situation for biological and economic systems, except that one may presuppose an underlying but unknown mathematical system description.

parameters of the driving noise. The information obtained then characterizes the situation fully. That is precisely what the paper does for the more general statistical model it considers. But the paper goes a step further. It does not restrict itself to the filtering situation.





Presenting Robert Stelzer's Primary Sources Essay

**Featured Essay** 

Foreword

**Author Profile** 

About Primary Sources

Often, the filter is just an input-output description of a more complex structure, in the system theory literature called a "dynamical system" - i.e., a system that contains a physical state, which evolves in time. The original Kalman filter is actually a "state estimator." Under Gaussian assumptions it makes sense to predict the state as well as possible on the basis of past observations; the difference between the predicted state and the actual one, called the "innovation," will be fully random and as small as possible. When the system is driven by a Lévy-Pareto process, the situation is very different. Least squares state prediction does not make much sense any more; much more important is to estimate the regime the system is in, as it can undergo drastic changes. Stelzer addresses the issue to estimate both the parameters of the system and the statistics of the driving process, hence obtaining maximal information compatible with the stochastic situation.

One of the goals of our publication, Primary Sources, is to inform a scientifically savvy audience about important new concepts and methods that may influence their thinking.

Number one in the series is the paper by Stelzer, and fittingly so, as it presents not only an important new technique, but also one with a wealth of potential applications. The **TUM-IAS Focus Group of Klüppelberg** and Stelzer has gone to great lengths connecting with potential appliers: designers of wind parks (for renewable energy), groups that are interested in deriving phenotypes from DNA, civil engineers studying risks associated with the structures they build, economists trying to minimize risk in trading, ecologists wanting to understand the statistical properties of natural processes, electrical engineers designing smart grids, and even fundamental physicists studying the erratic phenomena of nanosystems. Systems driven by Lévy-Pareto processes are indeed everywhere. And for new tools that could help "get a grip" on them all, scientists and engineers need look no further than these pages.

Professor Patrick Dewilde, director of the TUM Institute for Advanced Study and scientific editor of TUM-IAS Primary Souces, can be reached at dewilde@zv.tum.de.

#### TUM-IAS Primary Sources Featured Essay

Foreword

Featured Essay

**Author Profile** 

About Primary Sources



#### 1 Introduction

- 2 Lévy processes
- 3 Definition of CARMA processes and spectral representation
- **4** Properties
- 5 State space models
- 6 Statistical Estimation
- 7 Concluding Remarks

### CARMA processes driven by non-Gaussian noise

Robert Stelzer<sup>1</sup>

We present an outline of the theory of certain Lévy-driven, multivariate stochastic processes, where the processes are represented by rational transfer functions (Continuous-time AutoRegressive Moving Average or CARMA models) and their applications in non-Gaussian time series modeling. We discuss in detail their definition, their spectral representation, the equivalence to linear state space models and further properties like the second-order structure and the tail behavior under a heavy-tailed input. Furthermore, we study the estimation of the parameters using quasi-maximum likelihood estimates for the auto-regressive and moving average parameters, as well as how to estimate the driving Lévy process.

#### **1** Introduction

In many applications an observer (scientist, engineer, analyst) is confronted with series of data originating from one or more physical variables of interest over time. Thus, he or she has an observed (multivariate) time series and will often be interested either in removing (measurement) noise to extract the signal more clearly or in modeling the observed process, including its random components.

In both situations stochastic models may very well be appropriate. This is clear when one is mainly interested in removing noise; but it also is very often appropriate, when the intention is to model the observed value, to enrich a physical model by a random component to capture fluctuations and shortcomings of the physical model. The driving stochastic process (the "noise") may have interest on its own (as is the case with economic

Institute of Mathematical Finance, Ulm University, Helmholtzstraße 18, D-89069 Ulm, Germany. *Email:* robert.stelzer@uni-ulm.de, http://www.uni-ulm.de/mawi/finmath

Robert Stelzer: CARMA processes driven by non-Gaussian noise

models), or it may need to be modeled well to extract the interesting information as well as possible (e.g., as is common practice in telecommunication links).

The easiest way to obtain a model with randomness for the variables of interest would be to assume that all observed values are independent and identically distributed (iid) random variables or that they follow a physical model plus iid noise. However, in most series observed consecutive values are heavily dependent, and thus more sophisticated models are needed. A flexible but at the same time very tractable class of models is given by linear random processes. In the discrete time setting these models are well known as autoregressive moving average (ARMA) processes, and they are given in terms of a general order linear difference equation where an iid noisy input sequence introduces all randomness. The latter is also referred to as linear filtering of a white noise.

In many situations it is more appropriate to specify a model in continuous time rather than in discrete time. These include situations involving high-frequency data, irregularly spaced data, or missing observations, or where estimation and inference at various frequencies is to be carried out. Moreover, many physical models are formulated in continuous time, making such an approach more natural.

In the following we consider linear random processes in continuous time, referred to as continuous time autoregressive moving average (CARMA) processes. Intuitively, they are given as the solution to a higher-order system of linear differential equations with a stochastic process as the input, which can be seen as linearly filtering the random input.

One important question is which random input to take in the continuous time set-up. Clearly, the random process should correspond in some sense to the idea of white noise. Understanding the latter in the strict sense means using independent increments; in the weak sense it means uncorrelated increments, and so the variance has to be finite. Recall that for random variables uncorrelatedness is equivalent to independence only if the random variables are Gaussian, i.e., they have a normal distribution. A linear random process driven by Gaussian white noise has again Gaussian distributions. However, in many situations it is not appropriate to assume Gaussianity of the variables of interest, since the observed time series often exhibit features like skewness or heavy tails (i.e., very high or low values are far more likely to occur than in the Gaussian setting), which contradict the Gaussian assumption. Demanding uncorrelated but not necessarily

Featured Essay

**Author Profile** 

About Primary Sources

Robert Stelzer: CARMA processes driven by non-Gaussian noise

independent increments does not lead to a nice class of processes nor to nice theoretical results.

Hence, a good modeling strategy where the resulting process is reasonably tractable and the driving process's probability distribution is allowed to have "fat tails" is to demand that the random input must have independent as well as stationary increments; i.e., increments over time intervals of the same length have the same distribution. They then have a time homogeneity feature and resemble the iid noise of the discrete time set-up. The resulting class of possible driving processes are the so-called Lévy processes, which have been studied in detail and form a both highly versatile and highly tractable family. An interesting feature is that linear processes driven by general Lévy processes may exhibit jumps and thus allow the modeling of abrupt changes, whereas Gaussian linear processes have continuous sample paths.

In the remainder of this paper we proceed as follows. First, we introduce Lévy processes in detail. Thereafter, we give a proper definition of CARMA processes, discuss their relation to linear filtering via a stochastic Fourier (spectral) representation, and summarize central properties of CARMA processes. Next, we briefly explain the equivalence to linear state space models and the relation to stochastic control and signal processing. Finally, we discuss the statistical estimation of the parameters and the underlying Lévy process and conclude with some additional remarks.

Throughout we will focus on developing the main ideas for CARMA processes. For more mathematical details as well as comprehensive references we refer the interested reader to the original literature, especially the works [2], [8–10], [15–17], [12], [13], [42] and [55]. For a historic perspective the monograph [47] may be interesting as well as [19], which is the first paper where Gaussian CARMA processes appeared under the name of Gaussian processes with rational spectral density.

#### 2 Lévy processes

A Lévy process  $L = (L_t)_{t \in \mathbb{R}^+}$  is a stochastic process with independent and stationary increments. In the following we consider only Lévy processes taking values in the m-dimensional vector space  $\mathbb{R}^m$  (with  $\mathbb{R}$  the real numbers and m some positive integer). Note that a stochastic process  $(X_t)_{t \in \mathbb{R}^+}$  can be seen either as a family of random variables indexed by the

Featured Essay

Foreword

**Author Profile** 

About Primary Sources

**Featured Essay** 

**Author Profile** 

About Primary Sources

#### TUM-IAS Primary Sources

Robert Stelzer: CARMA processes driven by non-Gaussian noise

positive real numbers  $\mathbb{R}^+$  or as a random function mapping the positive real numbers to  $\mathbb{R}^m$ . More precisely we have the following definition:

**Definition 2.1.** An  $\mathbb{R}^m$ -valued stochastic process  $L = (L_t)_{t \in \mathbb{R}^+}$  is called Lévy process *if* 

- $L_0 = 0 \ a.s.,$
- $L_{t_2} L_{t_1}, L_{t_3} L_{t_2}, \dots, L_{t_n} L_{t_{n-1}}$  are independent for all  $n \in \mathbb{N}$  and  $t_1, t_2, \dots, t_n \in \mathbb{R}$ + with  $0 \leq t_1 < t_1 < \dots < t_n$ ,
- $L_{t+h} L_t \stackrel{\mathscr{D}}{=} L_{s+h} L_s$  for all  $s, t, h \in \mathbb{R}^+$  (" $\stackrel{\mathscr{D}}{=}$ " denoting equality in *distribution*),
- L is continuous in probability, i.e. for all  $s \in \mathbb{R}^+$  we have  $L_t L_s \xrightarrow{P} 0$  as  $t \to s$ .

It can be shown (cf. [53] for a detailed proof) that the class of Lévy processes can be characterized fully at the level of "characteristic functions," which we now introduce. Let  $\langle \cdot, \cdot \rangle$  indicate the natural inner product in  $\mathbb{R}^m$  and X is an  $\mathbb{R}^m$ -valued random variable; then its *characteristic function* is defined as  $\psi_X(u) = \mathbb{E}(e^{i \langle u, X \rangle})$ . The characteristic function of a Lévy process can always be represented in the *Lévy-Khintchine form* 

$$\mathsf{E}\left(e^{\mathsf{i}\langle \mathsf{u},\mathsf{L}_{\mathsf{t}}\rangle}\right) = \exp\{\mathsf{t}\psi_{\mathsf{L}}(\mathsf{u})\}, \ \forall \, \mathsf{t} \ge 0, \, \mathsf{u} \in \mathbb{R}^{\mathfrak{m}}, \tag{2.1}$$

with

$$\psi_{\mathrm{L}}(\mathfrak{u}) = \mathfrak{i}\langle\gamma,\mathfrak{u}\rangle - \frac{1}{2}\langle\mathfrak{u},\Sigma_{\mathrm{G}}\mathfrak{u}\rangle + \int_{\mathbb{R}^{m}} (e^{\mathfrak{i}\langle\mathfrak{u},\mathfrak{x}\rangle} - 1 - \mathfrak{i}\langle\mathfrak{u},\mathfrak{x}\rangle\mathbf{1}_{[0,1]}(\|\mathfrak{x}\|)\,\nu(d\mathfrak{x}),$$
(2.2)

where  $\gamma \in \mathbb{R}^m$ ,  $\Sigma_G$  is an  $m \times m$  positive semi-definite matrix and  $\nu$  is a measure on  $\mathbb{R}^m$  that satisfies  $\nu(\{0\}) = 0$  and  $\int_{\mathbb{R}^m} (\|x\|^2 \wedge 1) \nu(dx) < \infty$ . The measure  $\nu$  is referred to as the Lévy measure of L, and  $\|x\|^2 \wedge 1$  is short for  $\min\{\|x\|^2, 1\}$ . Finally,  $1_A(x)$  generically denotes the indicator function of a set A, i.e., the function that is one if x is an element of A and zero otherwise. Together  $(\gamma, \Sigma_G, \nu)$  are referred to as the characteristic triplet of L.

Robert Stelzer: CARMA processes driven by non-Gaussian noise

Regarding the paths of a Lévy process, i.e., the "curve of L as a function of time t," it can be shown that without loss of generality, a Lévy process may be assumed to be right continuous and have left limits.

It should be noted that many well known stochastic processes are Lévy processes. Examples are Brownian motion, also referred to as the Wiener process or "Gaussian white noise"; the Poisson process, which has jumps of size one and remains constant in between the jumps, which occur after iid exponentially distributed waiting times; and  $\alpha$ -stable Lévy motions, sometimes called Lévy flights. Compound Poisson processes are Poisson processes where the fixed jump size one is replaced by random iid jump sizes independent of the interarrival times of the jumps. It can be shown that all Lévy processes arise as limits of such compound Poisson processes.

A better understanding of what Lévy processes really are is provided by the Lévy-Itô decomposition of their paths. It states that a Lévy process is the sum of the deterministic linear function  $\gamma t$ , a Brownian motion with covariance matrix  $\Sigma_G$ , the sum of the big jumps that form a compound Poisson process, and the compensated sum of the small jumps (i.e., the sum of the small jumps minus their expected value). The quantity  $\nu(A)$  gives for any measurable set  $A \subset \mathbb{R}^m$  the expected number of jumps with size in A occurring in a time interval of length one. In Figure 1 a univariate Lévy process, which is the sum of the linear function t, in this case with  $\gamma = 2$ , a standard Brownian motion, with  $\Sigma_G = 1$ , and a Poisson process, with  $\nu(\{1\}) = 1$ ,  $\nu(\mathbb{R} \setminus \{1\}) = 0$  is depicted together with its individual components.

Whenever  $\int_{\mathbb{R}^m} (||x|| \wedge 1)\nu(dx) < \infty$ , we can simply replace the compensated sum of small jumps with the sum of the small jumps, adjusting also the slope of the deterministic component. We have actually already done this in Figure 1 where the resulting slope of the deterministic function is  $\gamma - \int_{\mathbb{R}} x\nu(dx) = 1$ . If  $\nu(\mathbb{R}) < \infty$ ; we have finitely many jumps in any bounded time interval, and the jumps form actually a compound Poisson process. Otherwise, we have infinitely, but countably many jumps in any bounded time interval. The reason we have in general a component referred to as "the compensated sum of the jumps" (i.e., it results from a certain limiting procedure; see e.g., [53]) is that in general the jumps are not summable. This is equivalent to the fact that the paths have infinite variation, like Brownian motion. Infinite variation intuitively means that the curve described by the stochastic process over finite time intervals has an infinite length. Clearly, this means that the fluctuations of the process over small time intervals are rather vivid.

Featured Essay

**Author Profile** 

**Primary Sources** 

About

Technische Universität München / Institute for Advanced Study





Figure 1. A Lévy process and its components: The complete Lévy process is depicted in the lower right display. In the upper row the deterministic drift component is depicted on the left and the standard Brownian motion component on the right. The left display in the middle row shows the standard (rate one) Poisson component and the right one the Brownian motion and the deterministic component added together. In the last row on the left the Brownian component plus the Poisson jumps are depicted.

Note that the scaling of the y-axis is different in the individual plots.

Robert Stelzer: CARMA processes driven by non-Gaussian noise

Author Profile

**Featured Essay** 

Foreword

About Primary Sources



Figure 2. Simulation of a Normal Inverse Gaussian (NIG) Lévy process.

In Figures 2 and 3 you can see simulations of different pure jump Lévy processes; i.e., in these cases  $\gamma = 0$  and  $\Sigma = 0$ . So there is neither a deterministic drift nor a Brownian motion present. All these processes have infinite activity, i.e., infinitely many jumps in any time interval. Figure 2 depicts a so-called normal inverse Gaussian Lévy process which has heavier tails than a Brownian motion, but still is rather tame, because it has finite moments of all orders, i.e.,  $E(|L_t|^r) < \infty$  for all  $t, r \in \mathbb{R}^+$ , and also some exponential moments. In contrast to this the stable processes of Figure 3 have very heavy tails, because they do not have a finite variance and the 0.5-stable process have infinite variation, the small jumps of the 0.5-stable Lévy process are summable.

Most of the time we will work with Lévy processes defined on the whole real line, i.e., indexed by  $\mathbb{R}$  not  $\mathbb{R}^+$ . They are obtained by taking two independent copies of a Lévy process and reflecting one copy at the origin.

For detailed expositions on Lévy processes we refer to [1], [6], [37] or [53].

1.5–Stable Process

Robert Stelzer: CARMA processes driven by non-Gaussian noise

#### Author Profile

Featured Essay

Foreword

About Primary Sources





Figure 3. Simulations of stable Lévy processes. A 1.5-stable Lévy process is depicted in the upper row and a 0.5-stable in the lower one.

**Featured Essay** 

**Author Profile** 

About Primary Sources Robert Stelzer: CARMA processes driven by non-Gaussian noise

## 3 Definition of CARMA processes and spectral representation

On the intuitive level one wants to be able to interpret a d-dimensional CARMA(p,q) process Y as the stationary solution to the p-th order linear differential equation

$$P(D)Y_{t} = (D^{p} + A_{1}D^{p-1} + \ldots + A_{p})Y_{t}$$
(3.1)

$$= (B_0 D^{\mathfrak{q}} + B_1 D^{\mathfrak{q}-1} + \ldots + B_{\mathfrak{q}}) DL_t = Q(D) DL_t, \qquad (3.2)$$

where the driving input L is an m-dimensional Lévy process, D denotes differentiation with respect to t, and the coefficients  $A_1, \ldots, A_p$  are  $d \times d$  matrices and  $B_0, \ldots, B_q$  are  $d \times m$  matrices. The polynomials  $P(z) = z^p + A_1 z^{p-1} + \ldots + A_p$  and  $Q(z) = B_0 z^q + B_1 z^{q-1} + \ldots + B_q$  with  $z \in \mathbb{C}$  are referred to as the auto-regressive and the moving average polynomial, respectively. Finally,  $p, q \in \mathbb{N}$  are the auto-regressive and moving average order.

However, the paths of non-deterministic Lévy processes are not differentiable, and so the above equation cannot directly provide a rigorous mathematical definition. Let us briefly consider the case (p, q) = (1, 0), in which case the resulting process is called an Ornstein-Uhlenbeck (OU) process. In the univariate case it is given by the differential equation

$$DY_t = aY_t + DL_t$$

where a is a real number. So what we basically want is that the change of Y over an infinitesimal time interval is a times the current value of the process times the "length of the infinitesimal time interval" plus the change of the Lévy process over the infinitesimal time interval. Rephrasing this idea in the precise language of stochastic differential equations (see e.g., [48]) we obtain

$$dY_t = aY_t dt + dL_t.$$

Using the theory of stochastic differential equations (SDEs), we can easily see that this SDE has a unique solution given by

$$Y_t = e^{at}Y_0 + e^{at}\int_0^t e^{-as}dL_s.$$

For general orders (p, q) one could to some extent use similar reasoning to arrive at a precise definition of CARMA processes. However, we shall take a more elegant route. First note that the differential operators on the

Robert Stelzer: CARMA processes driven by non-Gaussian noise

auto-regressive side of (3.1) act like integration operators on the moving average side. Hence, they offset the differential operators of the moving average side acting on the Lévy process. Since Lévy processes are not differentiable, we effectively have to integrate at least as often as we differentiate to be able to make sense of (3.1). Hence, a necessary condition ensuring the proper existence of CARMA processes is p > q.

In order to obtain a rigorous definition of CARMA processes our strategy here shall be to switch from the time domain to the frequency domain, where the main tool is the following spectral representation of a Lévy process. Here and in the following we denote by  $A^*$  for a matrix (or vector) A the Hermitian, i.e., the complex conjugate transposed matrix.

**Theorem 3.1** ([42]). Let  $(L_t)_{t\in\mathbb{R}}$  be a square integrable m-dimensional Lévy process with mean  $E[L_1] = 0$  (which implies  $E[L_t] = 0$  for all t) and variance  $E[L_1L_1^*] = \Sigma_L$  at t = 1. Then there exists a unique m-dimensional random orthogonal measure  $\Phi_L$  with spectral measure  $F_L$  such that  $E[\Phi_L(\Delta)] = 0$  for any bounded Borel set  $\Delta$ ,  $F_L(dt) = \frac{\Sigma_L}{2\pi} dt$  and

$$L_t = \int_{-\infty}^{\infty} \frac{e^{i\mu t} - 1}{i\mu} \Phi_L(d\mu), \ t \in \mathbb{R}.$$

*The random measure*  $\Phi_L$  *is uniquely determined by* 

$$\Phi_{\rm L}([a,b)) = \int_{-\infty}^{\infty} \frac{e^{-i\mu a} - e^{-i\mu b}}{2\pi i\mu} \, dL_{\mu} \tag{3.3}$$

*for all*  $-\infty < a < b < \infty$ .

The random orthogonal measure  $\Phi_L$  can intuitively be thought of as the "Fourier transform" of the Lévy process. If  $L_t$  is a Brownian motion, then  $\Phi_L([0, t))$  is again a Brownian motion. For general Lévy processes rather little can be said about the properties of  $\Phi_L$ . For example, it is known that  $\Phi_L$  has second-order stationary and uncorrelated increments, but the increments are neither independent nor stationary in a strict sense; see [27].

In the spectral domain we can now interpret differentiation (and integration) as linear filtering, noting that a formal interchange of differentiation and integration gives " $DL_t = \int_{-\infty}^{\infty} e^{i\mu t} \Phi_L(d\mu)$ ." It can be shown that the resulting process is well defined whenever the linear filter is square integrable. Thus we obtain as definition for "Y(t) = P(D)<sup>-1</sup>Q(D)DL(t)":

Foreword

Featured Essay

**Author Profile** 

About Primary Sources

Robert Stelzer: CARMA processes driven by non-Gaussian noise

**Definition 3.2** (CARMA Process, [42]). Let  $L = (L_t)_{t \in \mathbb{R}}$  be a two-sided square integrable m-dimensional Lévy-process with  $E[L_1] = 0$  and  $E[L_1L_1^*] = \Sigma_L$ . A d-dimensional Lévy-driven continuous time autoregressive moving average process  $(Y_t)_{t \in \mathbb{R}}$  of order (p, q) with  $p, q \in \mathbb{N}_0$  and p > q (CARMA(p, q) process) is defined as

$$\begin{split} Y_{t} &= \int_{-\infty}^{\infty} e^{i\mu t} P(i\mu)^{-1} Q(i\mu) \, \Phi_{L}(d\mu), \quad t \in \mathbb{R}, \qquad \textit{where (3.4)} \\ P(z) :&= I_{m} z^{p} + A_{1} z^{p-1} + \ldots + A_{p}, \\ Q(z) :&= B_{0} z^{q} + B_{1} z^{q-1} + \ldots + B_{q} \quad \textit{and} \end{split}$$

 $\Phi_{L}$  is the Lévy orthogonal random measure of Theorem 3.1. Here  $A_{j} \in M_{m}(\mathbb{R}), j = 1, ..., p$  and  $B_{j} \in M_{d,m}(\mathbb{R})$  are matrices satisfying  $B_{q} \neq 0$  and  $\mathcal{N}(P) := \{z \in \mathbb{C} : \det(P(z)) = 0\} \subset \mathbb{R} \setminus \{0\} + i\mathbb{R}$  (i.e. the autoregressive polynomial has no zeros on the complex axis).

Referring to the explicit construction of the random orthogonal measure  $\Phi_L$ , one can easily show that the above defined CARMA processes are necessarily stationary (in the strict sense, i.e., the distributions are left unchanged by a time shift). Since by construction any CARMA process in the sense of Definition 3.2 has a finite variance, it is also weakly stationary, i.e., the second-order moment structure (the variance and autocovariances) are left unchanged by time shifts.

Although the definition of CARMA processes via a spectral representation is elegant and helpful in many theoretical considerations, it is not really usable in applications, as alone simulating a CARMA process from this representation would be a tedious and problematic task. However, luckily we have the following result.

**Theorem 3.3** (State Space Representation, [42]). Let the Lévy process L and P, Q be as before. Define the following coefficient matrices:

• 
$$\beta_{p-j} = -\sum_{i=1}^{p-j-1} A_i \beta_{p-j-i} + B_{q-j}, \ j = 0, 1, \dots, q,$$
  
 $\beta_1 = \dots = \beta_{p-q-1} = 0$   
•  $\beta^* = (\beta_1^*, \beta_2^*, \dots, \beta_p^*) \text{ and } A = \left( \frac{0 | I_{d(p-1)}}{-A_p | -A_{p-1} - \dots -A_1} \right).$ 

**Featured Essay** 

Foreword

**Author Profile** 

About Primary Sources

**Featured Essay** 

**Author Profile** 

About Primary Sources Robert Stelzer: CARMA processes driven by non-Gaussian noise

Denote by  $G_t = (G_{1,t}^*, \ldots, G_{p,t}^*)^* a pd$ -dimensional process and assume that  $\mathcal{N}(P) := \{z \in \mathbb{C} : det(P(z)) = 0\} \subset (-\infty, 0) + i\mathbb{R}$  - the open right half of the complex plane. Then

$$dG_t = AG_t dt + \beta dL_t \tag{3.5}$$

has a unique stationary solution G given by

$$G_{t} = \int_{-\infty}^{t} e^{A(t-s)} \beta \, dL_{s}, \quad t \in \mathbb{R}.$$
(3.6)

It holds that

$$G_{1,t} = \int_{-\infty}^{\infty} e^{i\mu t} P(i\mu)^{-1} Q(i\mu) \Phi_{L}(d\mu) = Y_{t}, \ t \in \mathbb{R}.$$

So the first d-components of G are the CARMA process Y.

#### A CARMA process satisfying

 $\mathcal{N}(\mathsf{P}) := \{z \in \mathbb{C} : \det(\mathsf{P}(z)) = 0\} \subset (-\infty, 0) + i\mathbb{R} \text{ is called causal, because}$ as shown above the value at a time t only depends on the Lévy process up to time t; it is a function of  $(L_s)_{s \in (-\infty,t)}$ . In other words, a causal CARMA process is fully determined by values in the past. Whenever the condition  $\mathcal{N}(\mathsf{P}) := \{z \in \mathbb{C} : \det(\mathsf{P}(z)) = 0\} \subset (-\infty, 0) + i\mathbb{R} \text{ is not satisfied, } \mathsf{Y}_t \text{ also}$ depends on future values of the Lévy process. In many applications, where it is clear that all we see today can only be influenced by what happened up to now, one only considers causal processes as appropriate models. However, there are also applications where non-causal processes are useful. For example, if we want to stochastically model the water level in a river and think of t as describing the location along the river, both the water levels downstream (in the "future") and upstream (in the "past") may influence the water level at a certain point. Note that in this paper we only discuss stationary CARMA processes. In some applications (e.g., control) it is often adequate to consider non-stationary (non-stable) systems. Then the roots det(P(z)) = 0} in the set  $(-\infty, 0) + i\mathbb{R}$  describe the stable and causal part of the system, and the remaining roots describe the non-stable part.

Theorem 3.3 allows us to treat a causal CARMA process as a solution to the stochastic differential equation (3.5), and thus we can apply all the available results for SDEs. In particular, tasks like simulation of a causal CARMA process are straightforward and easily implemented. However, the above result allows us also to get rid of another restriction. So far we could only define CARMA processes driven by Lévy processes with finite second

**Featured Essay** 

**Author Profile** 

About Primary Sources TUM-IAS Primary Sources

Robert Stelzer: CARMA processes driven by non-Gaussian noise

moments, and thus we could so far not have, e.g., CARMA processes driven by  $\alpha$ -stable Lévy processes. However, general theory on multidimensional Ornstein-Uhlenbeck processes (see [36] and [54]) tells us that (3.6) is the unique stationary solution to (3.5) as soon as the Lévy process has only a finite logarithmic moment.

**Definition 3.4** (Causal CARMA Process, [42]). Let  $L = (L_t)_{t \in \mathbb{R}}$  be an *m*-dimensional Lévy process satisfying

$$\int_{\|x\|\ge 1} \ln \|x\| \,\nu(dx) < \infty, \tag{3.7}$$

p,  $q \in \mathbb{N}_0$  with q < p, and further  $A_1, A_2, \ldots, A_p, \in M_d(\mathbb{R})$ ,  $B_0, B_1, \ldots$ ,  $B_q \in M_{d,m}(\mathbb{R})$ , where  $B_0 \neq 0$ . Define the matrices A,  $\beta$  and the polynomial P as in Theorem 3.3 and assume  $\sigma(A) = \mathcal{N}(P) \subseteq (-\infty, 0) + i\mathbb{R}$ . Then the d-dimensional process

$$Y_t = (I_d, 0, \dots, 0) G_t$$
 (3.8)

where  $G_t = \int_{-\infty}^t e^{A(t-s)} \beta dL_s$  is the unique stationary solution to  $dG_t = AG_t dt + \beta dL_t$  is called causal CARMA(p, q) process.

G is referred to as the state space representation.

A natural question is clearly whether one can also extend the definition of CARMA processes via the spectral representation to the case with infinite variance. For so-called regularly varying Lévy processes with finite mean, and thus especially for  $\alpha$ -stable Lévy processes with  $\alpha \in (1, 2)$ , a result like Theorem 3.1 has been established in [27]. However, the non-finite variance case is distinctly different, as a limit of integrals has to be taken and the random orthogonal measure is replaced by an object that is – strictly speaking – not even a measure any more. In that paper a definition of CARMA processes with regularly varying Lévy input analogous to Definition 3.2 has been given, and it has been shown that the resulting processes coincide with the causal CARMA processes when both definitions apply. Observe that processes with infinite variance are not only of academic interest, but that they have important applications, for instance, in network data modeling (cf. [43] and [50, 51]). In [29] CARMA processes driven by  $\alpha$ -stable Lévy processes have been successfully used to model electricity prices.

**Featured Essay** 

**Author Profile** 

About Primary Sources **TUM-IAS Primary Sources** 

Robert Stelzer: CARMA processes driven by non-Gaussian noise

#### **4** Properties

In this section we explain and summarize various properties of (causal) CARMA processes.

#### 4.1 Second-Order Structure

Recall that for convenience we have assumed that the driving Lévy process (and thus the CARMA process) has mean zero. Looking at the "defining" differential equations, it is clear that if  $E(L_1) = \mu$  then the CARMA process is defined as the one driven by  $L_t - \mu t$  plus  $A_p^{-1}B_q\mu$ , which is then the mean of the CARMA process.

**Proposition 4.1** ([42]). *Let* Y *be a (causal) CARMA process driven by a Lévy process* L *with finite second moments and set*  $\Sigma_L = var(L_1)$ .

1. The CARMA process Y has autocovariance function:

$$\operatorname{cov}(Y_{t+h},Y_t) = \int_{-\infty}^{\infty} \frac{e^{i\mu h}}{2\pi} P(i\mu)^{-1} Q(i\mu) \Sigma_L Q(i\mu)^* (P(i\mu)^{-1})^* \, d\mu,$$

with  $h \in \mathbb{R}$ .

2. If Y is a causal CARMA process, its state space representation G has the following second-order structure:

$$\begin{aligned} \text{var}(G_t) &= \int_{0}^{\infty} e^{Au} \beta \Sigma_L \beta^* e^{A^*u} du \\ \text{Avar}(G_t) + \text{var}(G_t) A^* &= -\beta \Sigma_L \beta^* \\ \text{cov}(G_{t+h}, G_t) &= e^{Ah} \text{var}(G_t), \ h \ge 0. \end{aligned}$$

Since we are only considering stationary CARMA processes, the moments above do not depend on t.

Since Y is given by the first d components of G, the second-order structure of G implies immediately alternative formulae for the second-order structure of Y. In particular, it shows that the autocovariance function always decays like a matrix exponential for  $h \rightarrow \infty$ .

**Featured Essay** 

**Author Profile** 

About Primary Sources Robert Stelzer: CARMA processes driven by non-Gaussian noise

#### 4.2 Distribution

Another nice feature is that in principle the distribution of a CARMA process at fixed times and the higher-dimensional marginal distributions, e.g., the joint distribution of the process at two (or n) different points in time, is explicitly known in terms of the characteristic function. The reasons are that all these distributions are infinitely divisible and that their Lévy-Khintchine triplet is known in terms of the Lévy-Khintchine triplet of the driving Lévy process. We state this in detail for the stationary distribution in the causal case.

**Proposition 4.2** ([42]). If L has characteristic triplet  $(\gamma, \Sigma, \nu)$ , then the stationary distribution of the state space representation G of a causal *CARMA* process is infinitely divisible with characteristic triplet  $(\gamma_{G}^{\infty}, \Sigma_{G}^{\infty}, \nu_{G}^{\infty})$ , where

• 
$$\gamma_{G}^{\infty} = \int_{0}^{\infty} e^{As} \beta \gamma \, ds + \int_{0}^{\infty} \int_{\mathbb{R}^{m}} e^{As} \beta x [1_{[0,1]}(\|e^{As} \beta x\|) - 1_{[0,1]}(\|x\|)] \nu(dx) \, ds,$$
  
•  $\Sigma_{G}^{\infty} = \int_{0}^{\infty} e^{As} \beta \Sigma \beta^{*} e^{A^{*}s} \, ds,$   
•  $\nu_{G}^{\infty}(B) = \int_{0}^{\infty} \int_{\mathbb{R}^{m}} 1_{B}(e^{As} \beta x) \nu(dx) \, ds$ 

for all Borel sets  $B \subseteq \mathbb{R}^{pd}$ .

In other words

$$E\left(e^{i\langle u,G_{t}\rangle}\right) = \exp\left\{i\langle\gamma_{G}^{\infty},u\rangle - \frac{1}{2}\langle u,\Sigma_{G}^{\infty}u\rangle + \int_{\mathbb{R}^{p\,d}} (e^{i\langle u,x\rangle} - 1 - i\langle u,x\rangle \mathbf{1}_{[0,1]}(\|x\|) \,\nu_{G}^{\infty}(dx)\right\}, \quad (4.1)$$

for all  $\mathfrak{u} \in \mathbb{R}^{pd}$ .

Projection onto the first d coordinates gives the characteristic triplet of the stationary distribution of Y. It should, however, be noted that typically the distribution of the CARMA process does not belong to any special family of distributions even if one starts with especially nice Lévy processes.

#### 4.3 Dependence Structure

An important property of multivariate stochastic processes is how their future evolution depends on the past. Suppose that one stands at a given

Featured Essay

Foreword

**Author Profile** 

About Primary Sources Robert Stelzer: CARMA processes driven by non-Gaussian noise

point in time and disposes of sufficient data at that point to determine the evolution from that point on, also given knowledge of the input from that point onwards. A *Markov process* is a stochastic process for which the future only depends on the current value and not any more on the past values (all their information is subsumed in the current value). For a Markov process it only matters – so to speak – where we are now, not where we came from. If this characterizing property does not only hold at all fixed times, but also at certain random times called stopping times, we speak of a strong Markov process.

**Proposition 4.3** ([42]). *The state space representation* G *of a causal CARMA process is a strong Markov process.* 

Intuitively it is desirable in many applications that the farther away observations are in time, the less dependent they should be. Usually, one even wants that very far away observations should be basically independent. This idea is mathematically formalized in various concepts of asymptotic independence often referred to as some form of "mixing."

A comparably weak result which, however, applies to any CARMA process is the following.

**Proposition 4.4** ([28]). Any stationary CARMA process is mixing.

Mixing implies ergodicity, i.e., empirically determined moments from the time series converge to the true moments if more and more data are collected. So time averages converge to ensemble averages. This is very important for statistical estimation of CARMA processes, as it implies typically that estimators are consistent (i.e., the estimators converge to the correct value when more and more data are collected).

Typically, one also wants to know the errors of estimators, which can be derived from distributional limit results like asymptotic normality. To obtain such results, a stronger, more uniform notion of asymptotic independence is needed, which is called strong mixing. Typically, one can best establish it for a Markov process.

**Proposition 4.5** ([42]). For a causal CARMA process with  $E(||L_1||^r) < \infty$  for some r > 0 the state space representation G and the CARMA process Y are strongly mixing, both with exponentially decaying mixing coefficients.

**Featured Essay** 

**Author Profile** 

About Primary Sources Robert Stelzer: CARMA processes driven by non-Gaussian noise

#### 4.4 Sample Path Properties

Next we look at the sample path properties of a CARMA process.

**Proposition 4.6** ([42]).

• The sample paths of a CARMA(p, q) process Y with p > q + 1 are (p - q - 1)-times differentiable, and for a causal CARMA process it holds that

$$\frac{d^{\iota}}{dt^{\iota}}Y_t = G_{\iota+1,t}, \quad i = 1, 2, \dots, p-q-1.$$

- If p = q + 1, and the driving Lévy process has a non-zero Lévy measure  $\nu$  satisfying  $\nu(B_0^{-1}(\mathbb{R}^d \setminus \{0\})) \neq 0$ , then the paths of a CARMA process exhibit jumps, and the jumps sizes are given by  $\Delta Y_t := Y_t - Y_{t-} = B_0 \Delta L_t$ .
- If the driving Lévy process L is a Brownian motion, then the sample paths of Y are continuous and (p − q − 1)-times continuously differentiable, provided p > q + 1.

For examples of the paths of CARMA processes driven by an NIG Lévy process see Figure 4.

#### 4.5 Tail behavior

As already stated in the introduction, one may want to move away from Gaussian models because extreme (i.e., very low and/or high) observations are far more likely than in a Gaussian distribution. One says that the tails (of the distribution) are heavier than Gaussian ones. Very often it appears also reasonable to use models that are "heavy-tailed" in the sense that only a limited number of moments exist, i.e.,  $E(||X||^r)$  exists only for low values of r. Mathematically it is then convenient to use the concept of regular variation (see [22] or [49, 51] for comprehensive introductions in relation to extreme value theory). Roughly speaking this means that the tails behave like a power function when one is far from the center of the distribution. A random variable X is regularly varying if P(||X|| > x) behaves comparably to  $x^{-\alpha}$  for some  $\alpha > 0$  and big values of x. In [44] (see also [23] in the univariate case) it is shown that under a very mild non-degeneracy condition a CARMA process driven by a regularly varying Lévy process is

Robert Stelzer: CARMA processes driven by non-Gaussian noise

Foreword

Featured Essay

**Author Profile** 

About Primary Sources



Figure 4. A CARMA(1,0) process driven by an NIG Lévy process having discontinuous paths is shown in the upper display and a CARMA(2,0) process driven by the same Lévy process having continuous paths in the lower one.

Robert Stelzer: CARMA processes driven by non-Gaussian noise

again regularly varying with the same index  $\alpha$ . Hence, it is straightforward to construct heavy-tailed CARMA processes when applications call for such features.

In the univariate case the tail behavior of CARMA processes is also understood in certain non-Gaussian situations, where one has lighter tails than regularly varying ones (see [24, 25]).

#### 5 State space models

We have defined the causal CARMA process using a so-called state space representation, and we have noted that the state space representation G is made up of the CARMA process Y and its derivatives as long as they exist. Hence, causal CARMA processes may be viewed as special state space models driven by Lévy processes. In fact, any state space model can also be realized as a CARMA process, as will be shown now.

We start with a precise definition of state space models.

Definition 5.1. Let L be an m-dimensional Lévy process and

 $A\in M_N(\mathbb{R}), \quad B\in M_{N,m}(\mathbb{R}), \quad C\in M_{d,N}(\mathbb{R}).$ 

A general (N, d)-dimensional continuous time state space model driven by L with parameters A, B, C is a solution of

 $\label{eq:thestate} \begin{array}{ll} \mbox{the state equation} & d\mathbf{X}_t = \!\!A\mathbf{X}_t dt + B d\mathbf{L}_t \\ \mbox{and the observation equation} & \mathbf{Y}_t = \!\!C\mathbf{X}_t. \end{array}$ 

X is called the state process and Y the output process.

Note that the state process is N-dimensional, whereas the output process is d-dimensional.

Sufficient conditions for the existence of a unique causal stationary solution of the state equation are given by  $(\mathfrak{R}(\cdot))$  indicates the "real part" of a complex number or function)

 $\Re(\lambda_{\nu}) < 0, \quad \lambda_{\nu}, \nu = 1, \dots, N$ , being the eigenvalues of A

and L having finite second moments.

Foreword

**Featured Essay** 

**Author Profile** 

**Primary Sources** 

About

**Featured Essay** 

**Author Profile** 

About Primary Sources TUM-IAS Primary Sources

Robert Stelzer: CARMA processes driven by non-Gaussian noise

It can easily be shown by integration that **X** satisfies

$$\mathbf{X}_{t} = e^{A(t-s)}\mathbf{X}_{s} + \int_{s}^{t} e^{A(t-u)} B d\mathbf{L}_{u}.$$

Likewise, the stationary output process Y satisfies

$$\mathbf{Y}_{t} = \int_{-\infty}^{t} C e^{A(t-u)} B d\mathbf{L}_{u}.$$

Its spectral density, the Fourier transform of the autocovariance function, is given by

$$f_{\mathbf{Y}}(\boldsymbol{\omega}) = \frac{1}{2\pi} C(i\boldsymbol{\omega} - A)^{-1} B \boldsymbol{\Sigma}_{\mathbf{L}} B^{\mathsf{T}} (-i\boldsymbol{\omega} - A^{\mathsf{T}})^{-1} C^{\mathsf{T}}.$$

From Definition 3.4 it is obvious that a CARMA process is a (pd, d)-dimensional state space model driven by an m-dimensional Lévy process. The following theorem states that also the converse is true.

**Theorem 5.2** ([55]). The stationary solution **Y** of the multivariate state space model (A, B, C, L) is an L-driven CARMA process with autoregressive polynomial P and moving average polynomial Q if and only if

$$C(zI_N - A)^{-1}B = P(z)^{-1}Q(z), \quad \forall z \in \mathbb{C}.$$

For any (A, B, C) there exist P, Q such that the above equation is satisfied and vice versa.

In reality we typically do not observe some variables of interest continuously, but only at a discrete set of points in time. Let us assume that we sample the process at an equidistant time grid with grid length h > 0 and denote by  $\mathbf{Y}_n^{(h)} := \mathbf{Y}_{nh}$  for  $n \in \mathbb{Z}$  the sampled observations of a state space process.

It is easy to see that

$$\mathbf{Y}_{n}^{(h)} = C\mathbf{X}_{n}^{(h)} \tag{5.1}$$

$$\mathbf{X}_{n}^{(h)} = e^{Ah} \mathbf{X}_{n}^{(h)} + \int_{(n-1)h}^{nn} e^{A(nh-u)} B d\mathbf{L}_{u},$$
(5.2)

which immediately shows that  $\mathbf{Y}_{n}^{(h)}$  is the output process of a discrete time (N, d)-dimensional state space model driven by the N-dimensional iid noise  $\left(\int_{(n-1)h}^{nh} e^{A(nh-u)} B d\mathbf{L}_{u}\right)_{n \in \mathbb{Z}}$ .

Robert Stelzer: CARMA processes driven by non-Gaussian noise

It is well known that any (N, d)-dimensional state space model in discrete time is an ARMA process. Combining this with Theorem 5.2 tells us that any equidistantly sampled CARMA process  $Y^{(h)}$  is an ARMA process. This observation will be the basis for estimating CARMA parameters in the next section, where we will need a considerable refinement of this result.

In many applications the sampling frequency is quite high, i.e., h is very small. Thus it is important to understand how  $Y^{(h)}$  behaves as  $h \to 0$ , which has been investigated in [18].

As we only observe the process **Y** in a state space model, an important question is what can be said about the state process **X** based on the observations. Hence, we want to reconstruct or "estimate" the latent process **X** as well as possible. This procedure is also referred to as filtering. For Gaussian state space models the easily implementable Kalman filter (see e.g. [11]) is optimal with respect to both variance and distribution. For non-Gaussian state space models with finite variance the very same procedure, now typically called linear filtering, gives an "estimate" of the latent process, which is the linear "estimate" (in the observations) with the lowest variance. However, it is typically not the "estimate" with the minimal variance and not a conditional expectation. Thus, more involved filtering techniques, such as particle filtering (see e.g. [20]), are better.

State space models, mainly Gaussian ones, are also heavily used in stochastic control (see [30] and references therein for a comprehensive overview) and signal processing (see [38, 39, 45], for instance). In both areas one is sometimes dealing with data for which a Gaussianity assumption is not really appropriate due to skewedness, excess kurtosis or heavy-tailedness. Clearly, in such situations Lévy-driven state space models or equivalently CARMA processes should be appealing. It seems worthwhile to mention that there are two uses of state space models in control, even though going into the details would be beyond the scope of this paper. Sometimes one assumes some random input that is then "controlled" by the state space model, so the the state space model acts as the controller. In contrast to this, sometimes the output of the state space model is regarded as the natural output of some system on which an additional controller is acting to ensure that the output meets certain requirements.

Foreword

Featured Essay

**Author Profile** 

About Primary Sources

**Featured Essay** 

**Author Profile** 

About Primary Sources Robert Stelzer: CARMA processes driven by non-Gaussian noise

#### **6** Statistical Estimation

In this section we discuss ways to estimate the parameters of a CARMA process and its driving Lévy process. First we address the estimation of the autoregressive and moving average parameters. Due to parametrization issues explained later on, we formally do this for Lévy-driven continuous time state space models, as defined in the previous section. In the univariate case quasi-maximum likelihood estimation of CARMA processes is comprehensively studied in [17].

#### 6.1 Quasi-maximum likelihood estimation

We assume that we observe the process Y at discrete, equally spaced times

$$\mathbf{Y}_{n}^{(h)} \coloneqq \mathbf{Y}_{nh}, \quad n \in \mathbb{Z}, \quad h > 0.$$

Furthermore, we define the linear innovations  $\varepsilon^{(h)}$  by

$$\boldsymbol{\varepsilon}_{n}^{(h)} = \boldsymbol{Y}_{n}^{(h)} - \boldsymbol{P}_{n-1} \boldsymbol{Y}_{n}^{(h)},$$

where  $P_{n-1}$  denotes the orthogonal projection onto  $\overline{\text{span}} \left\{ \mathbf{Y}_{\nu}^{(h)} : -\infty < \nu < n \right\}$ , i.e., the linear space spanned by the observations until time (n-1)h. From the construction it is immediate that  $(\boldsymbol{\epsilon}_{n}^{(h)})_{n \in \mathbb{Z}}$  is a white noise sequence, i.e., it has mean zero, a constant variance and is uncorrelated.

**Theorem 6.1** ([55]). Assume the eigenvalues  $\lambda_1, \ldots, \lambda_N$  of the matrix A are pairwise distinct and define complex numbers  $\Phi_1, \Phi_2, \ldots, \Phi_N$  by

$$1 - \Phi_1 z - \Phi_2 z^2 - \ldots - \Phi_N z^N = \prod_{\nu=1}^N \left[ 1 - e^{-\lambda_\nu h} z \right] \quad \forall z \in \mathbb{C}.$$

Then there exist  $\Theta_1, \Theta_2, \dots, \Theta_{N-1}$  in  $M_d(\mathbb{C})$  such that

$$\mathbf{Y}_{n}^{(h)} - \Phi_{1} \mathbf{Y}_{n-1}^{(h)} - \ldots - \Phi_{N} \mathbf{Y}_{n-N}^{(h)} = \varepsilon_{n}^{(h)} + \Theta_{1} \varepsilon_{n-1}^{(h)} + \ldots + \Theta_{N-1} \varepsilon_{n-N+1}^{(h)}$$

holds.

*Hence*,  $\mathbf{Y}^{(h)}$  *is a weak* ARMA(N, N - 1) *process.* 

This result suggests that one could estimate simply the ARMA coefficients of the sampled process and then transfer these estimates to estimates of the

**Featured Essay** 

Foreword

**Author Profile** 

About Primary Sources Robert Stelzer: CARMA processes driven by non-Gaussian noise

CARMA coefficients. However, to estimate a CARMA process it is not sufficient to estimate an ARMA process, because not all ARMA processes can be embedded in a CARMA process. There are ARMA processes that cannot arise as equidistantly sampled CARMA processes. The way out is to carry out the "ARMA estimation" in the CARMA parameter space.

Since we are going to use a quasi-maximum likelihood approach and have discretely sampled observations, all possible models considered in the estimation have to be distinguishable based only on the second-order properties of the sampled process.

**Definition 6.2** (Identifiability). A collection of continuous time stochastic processes  $(\mathbf{Y}_{\vartheta}, \vartheta \in \Theta)$  is identifiable if for any  $\vartheta_1 \neq \vartheta_2$  the two processes  $\mathbf{Y}_{\vartheta_1}$  and  $\mathbf{Y}_{\vartheta_2}$  have different spectral densities.

It is h-identifiable, h > 0, if for any  $\vartheta_1 \neq \vartheta_2$  the two processes  $\mathbf{Y}_{\vartheta_1}^{(h)}$  and  $\mathbf{Y}_{\vartheta_2}^{(h)}$  have different spectral densities.

We assume that our parametrization is given by a compact parameter space  $\Theta \subset \mathbb{R}^q$  with some  $q \in \mathbb{N}$  and a mapping

$$\psi: \Theta \ni \vartheta \mapsto (A_{\vartheta}, B_{\vartheta}, C_{\vartheta}, L_{\vartheta}).$$

We need to ensure that our parametrization is minimal regarding the dimensions, since a fixed output process can result from artificially arbitrarily high-dimensional state space models.

**Assumption P1** (Minimality). For all  $\vartheta \in \Theta$  the triple  $(A_{\vartheta}, B_{\vartheta}, C_{\vartheta})$  is minimal in the sense that if

$$C(zI_{\mathfrak{m}}-A)^{-1}B = C_{\vartheta}(zI_{N}-A_{\vartheta})^{-1}B_{\vartheta}$$

then  $m \ge N$  must be true.

**Assumption P2** (Eigenvalues). For all  $\vartheta \in \Theta$  the eigenvalues of  $A_{\vartheta}$  are pairwise distinct and contained in the strip

$$\{z \in \mathbb{C} : -\pi/h < \Im(z) < \pi/h\}.$$

We want to use a parametrization for the continuous time state space model, but need to ensure that it is h-identifiable. The following theorem provides easy-to-check criteria.

Featured Essay

**Author Profile** 

About Primary Sources

#### TUM-IAS Primary Sources

Robert Stelzer: CARMA processes driven by non-Gaussian noise

**Theorem 6.3** ([56]). Assume that the parameterization  $\psi : \Theta \supset \vartheta \mapsto (A_{\vartheta}, B_{\vartheta}, C_{\vartheta}, L_{\vartheta})$  is

- identifiable
- minimal
- and satisfies the eigenvalue condition.

Then the corresponding collection of output processes  $\{\mathbf{Y}_{\vartheta}, \vartheta \in \Theta\}$  is h-identifiable.

The quasi-maximum likelihood (QML) estimator is now obtained by pretending the observations were Gaussian, taking the corresponding likelihood, and maximizing it. More precisely the QML of  $\vartheta$  based on L observations  $\mathbf{y}^{L} = (\mathbf{y}_{1}, \dots, \mathbf{y}_{L})$  is

$$\hat{\boldsymbol{\vartheta}}^{\mathsf{L}} = \operatorname{argmax}_{\boldsymbol{\vartheta}\in\Theta} \mathcal{L}_{\boldsymbol{\vartheta}} \left( \mathbf{y}^{\mathsf{L}} \right),$$

where  $\mathcal{L}_{\vartheta}$  is the Gaussian likelihood function, which is proportional to

$$\left(\prod_{n=1}^{L} \det V_{\vartheta,n}\right)^{-1/2} \exp\left\{-\frac{1}{2} \sum_{n=1}^{L} \mathbf{e}_{\vartheta,n}^{\mathsf{T}} V_{\vartheta,n}^{-1} \mathbf{e}_{\vartheta,n}\right\}$$

with

$$\begin{split} \mathbf{e}_{\vartheta,n} = & \mathbf{y}_n - \mathsf{P}_{n-1} \mathbf{Y}_{\vartheta,n}^{(n)} \Big|_{\left\{\mathbf{Y}_{\vartheta,\nu}^{(h)} = \mathbf{y}_{\nu}: 1 \leq \nu < n\right\}}, \\ & \mathcal{V}_{\vartheta,n} = & \mathbb{E}\left[ \mathbf{e}_{\vartheta,n} \mathbf{e}_{\vartheta,n}^{\mathsf{T}} \left| \mathbf{Y}_{\vartheta,\nu}^{(h)} = \mathbf{y}_{\nu}: 1 \leq \nu < n \right. \right] \end{split}$$

So  $\mathbf{e}_{\vartheta,n}$  are the linear innovations under the model given by  $\vartheta$ , and  $V_{\vartheta,n}$  are their variances or the one-step prediction errors.

Computing the QML estimator is now a straightforward task utilizing the Kalman recursions and numerically maximizing the likelihood. However, since we have not used the true likelihood, it is not clear whether the resulting estimators are really sensible in the sense that they converge to the true parameters. Luckily, one can show that the estimators are well behaved.

**Theorem 6.4** (Strong consistency, [56]). For every sampling interval h > 0, the QML estimator  $\hat{\vartheta}^{L}$  is strongly consistent, i.e.

$$\hat{\boldsymbol{\vartheta}}^{\mathsf{L}} \rightarrow \boldsymbol{\vartheta}_{0} \quad a.s. \ as \ \mathsf{L} \rightarrow \infty,$$

provided the parametrization is h-identifiable.

**Featured Essay** 

**Author Profile** 

About Primary Sources

#### TUM-IAS Primary Sources

Robert Stelzer: CARMA processes driven by non-Gaussian noise

However, so far we cannot assess the quality of our estimators by confidence intervals, etc., which is made possible by the following result.

**Theorem 6.5** (Asymptotic normality, [56]). Assume that the driving Lévy process satisfies  $\mathbb{E}||\mathbf{L}(1)||^{4+\delta} < \infty$  for some  $\delta > 0$ . For every sampling interval h > 0, the QML estimator  $\hat{\vartheta}^{L}$  is asymptotically normally distributed, i.e.

$$\sqrt{\mathrm{L}}\left(\widehat{\boldsymbol{\vartheta}}^{\mathrm{L}}-\boldsymbol{\vartheta}_{0}\right) \xrightarrow{\mathcal{D}} \mathcal{N}(\boldsymbol{0},\Omega), \quad \Omega = \mathrm{J}(\boldsymbol{\vartheta}_{0})^{-1}\mathrm{I}(\boldsymbol{\vartheta}_{0})\mathrm{J}(\boldsymbol{\vartheta}_{0})^{-1},$$

with

$$\begin{split} J(\boldsymbol{\vartheta}) &= -\lim_{L \to \infty} \frac{2}{L} \frac{\partial^2}{\partial \boldsymbol{\vartheta} \partial \boldsymbol{\vartheta}^{\mathsf{T}}} \ln \mathcal{L}_{\boldsymbol{\vartheta}} \left( \mathbf{y}^{\mathsf{L}} \right), \\ I(\boldsymbol{\vartheta}) &= \lim_{L \to \infty} \frac{4}{L^2} \operatorname{Var} \frac{\partial}{\partial \boldsymbol{\vartheta}} \ln \mathcal{L}_{\boldsymbol{\vartheta}} \left( \mathbf{y}^{\mathsf{L}} \right), \end{split}$$

provided the parametrization is h-identifiable.

To obtain identifiable parametrizations one uses as in the discrete time case (see [32] or [40], for instance) so-called canonical parametrizations such as the echelon state space form. For more details on this we refer to [56]. Since such parametrizations are typically available for state space models rather than CARMA processes, one normally estimates state space models rather than the equivalent CARMA processes.

Let us finally look at one simulation study.

A d-dimensional normal inverse Gaussian (NIG) Lévy process L (see e.g. [3, 7, 46]) with parameters

$$\delta > 0, \kappa > 0, \beta \in \mathbb{R}^{d}, \Delta \in \mathsf{M}^{+}_{d}(\mathbb{R})$$

is given by a normal mean-variance mixture, i.e.

$$\mathbf{L}_1 = \boldsymbol{\mu} + \mathbf{V} \Delta \boldsymbol{\beta} + \mathbf{V}^{1/2} \mathbf{N},$$

where N is d-dimensionally normally distributed with mean zero and variance  $\Delta$  and independent of

$$V \sim IG(\delta/\kappa, \delta^2)$$

which follows a so-called inverse Gaussian distribution ([35]).

We consider now a bivariate NIG-driven CARMA process with zero mean given by the state space form

**Featured Essay** 

**Author Profile** 

About Primary Sources

$$\begin{split} d\mathbf{X}_{t} &= \begin{bmatrix} \vartheta_{1} & \vartheta_{2} & 0 \\ 0 & 0 & 1 \\ \vartheta_{3} & \vartheta_{4} & \vartheta_{5} \end{bmatrix} \mathbf{X}_{t} dt + \begin{bmatrix} \vartheta_{1} & \vartheta_{2} \\ \vartheta_{6} & \vartheta_{7} \\ \vartheta_{3} + \vartheta_{5} \vartheta_{6} & \vartheta_{4} + \vartheta_{5} \vartheta_{7} \end{bmatrix} d\mathbf{L}_{t}, \\ \mathbf{Y}_{t} &= \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix} \mathbf{X}_{t}, \quad \boldsymbol{\Sigma}_{L} = \begin{bmatrix} \vartheta_{8} & \vartheta_{9} \\ \vartheta_{9} & \vartheta_{10} \end{bmatrix}. \end{split}$$

Robert Stelzer: CARMA processes driven by non-Gaussian noise

The parameters are  $\vartheta_1, \vartheta_2, \ldots, \vartheta_{10}$ , and the parametrization is in one of the canonical identifiable forms.

A simulated path is shown in Figure 5.

We calculated the QML estimates for this bivariate NIG-driven CARMA process on the basis of observations over the time horizon [0, 2000] at integer times and repeated this for 350 different simulated paths. The estimation results are summarized in Table 1. It shows that the sample bias of the obtained estimators in the simulation study is very small and that the sample standard deviation is close to the standard deviation predicted by the asymptotic normality result Theorem 6.5. Actually, the sample standard deviation is always smaller, which is nice as it implies that the standard deviation predicted by the asymptotic normality result Theorem 6.5 is a conservative estimate.

#### 6.2 Statistical inference for the driving Lévy process

The above quasi-maximum likelihood approach only allows one to estimate the autoregressive and moving average parameters and the variance of the driving Lévy process. However, typically we want to estimate many more parameters of the driving Lévy process or even first need to get an idea which family the driving Lévy process may belong to. To this end, one can reconstruct from the CARMA process the driving Lévy process. Typically, the CARMA process is only observed at a discrete set of times, and then the best we can do is to get approximations of the increments of the Lévy process. One can then treat the approximate increments as if they were the true ones of the Lévy process. "Looking" at them, one should be able to choose appropriate parametric families. By using the approximate increments as one would use the true ones (in maximum likelihood or method of moment-based estimation procedures), one can do parametric inference for the Lévy process. The construction of the approximate

Featured Essay

**Author Profile** 

About Primary Sources



Robert Stelzer: CARMA processes driven by non-Gaussian noise

Figure 5. One realization of a bivariate NIG-driven CARMA process (upper two displays) and the effect of sampling (lower two displays). The linearly interpolated process over the time interval [600, 650] resulting from sampling at integer times is shown as the thicker line, whereas the thinner line is the true CARMA process.

increments and their use in estimation procedures have been studied in detail in [14], where it is shown in particular that the estimators are good in the sense that they are consistent and asymptotically normal under reasonable assumptions when taking appropriate limits.

It should be noted that the idea to reconstruct the Lévy process can already be found in [10] or [17]. In the following we illustrate this approach for a univariate Ornstein-Uhlenbeck, i.e., a CARMA(1,0) process based on [16] and [31], from which all examples and plots are taken.

**Featured Essay** 

**Author Profile** 

About Primary Sources Robert Stelzer: CARMA processes driven by non-Gaussian noise

Table 1. Summary of the results of the simulation study on the QML estimation of a bivariate NIG-driven CARMA process. The second column states the mean of estimators obtained over 350 simulated paths, the third column the resulting bias, and the fourth column the standard deviation of the obtained estimators. Finally, the last column states the standard deviation for the estimators as predicted by the asymptotic normality result Theorem 6.5.

parameter	sample mean	sample bias	sample standard deviation	estimated standard deviation
$\vartheta_1$	-1.0001	0.0001	0.0354	0.0381
$\vartheta_2$	-2.0078	0.0078	0.0479	0.0539
$\vartheta_3$	1.0051	-0.0051	0.1276	0.1321
$\vartheta_4$	-2.0068	0.0068	0.1009	0.1202
$\vartheta_5$	-2.9988	-0.0012	0.1587	0.1820
$\vartheta_6$	1.0255	-0.0255	0.1285	0.1382
$\vartheta_7$	2.0023	-0.0023	0.0987	0.1061
$\vartheta_8$	0.4723	-0.0028	0.0457	0.0517
θ9	-0.1654	0.0032	0.0306	0.0346
$\vartheta_{10}$	0.3732	0.0024	0.0286	0.0378

Recall that an Ornstein-Uhlenbeck (OU) process is the unique strictly stationary solution to

$$dY_t = aY_t dt + dL_t. (6.1)$$

where  $(L_t)_{t\in\mathbb{R}}$  is a Lévy process with  $E(\ln(max(|L_1|, 1))) < \infty$  and autoregressive parameter a < 0. The solution of the stochastic differential equation is given explicitly by

$$Y_{t} = e^{a(t-s)}Y_{s} + \int_{s}^{t} e^{a(t-u)}dL_{u}.$$
(6.2)

If the OU process is observed continuously on [0, T], then the integrated form of (6.1) immediately gives

$$L_t = Y_t - Y_0 - \alpha \int_0^t Y_s ds.$$

The increments of the driving Lévy process  $\Delta L_n^{(h)}$  on the intervals ((n-1)h, nh] with  $n \in \mathbb{N}$  can be represented as

$$\Delta L_{n}^{(h)} := L_{nh} - L_{(n-1)h} = Y_{nh} - Y_{(n-1)h} - a \int_{(n-1)h}^{nn} Y_{u} du.$$
 (6.3)

Robert Stelzer: CARMA processes driven by non-Gaussian noise

What we want is to approximately reconstruct the sequence  $\Delta L^{(h)}$  of increments over intervals of length h from observations of the CARMA process made over a finer equidistant grid. To this end one simply approximates the integral  $\int_{(n-1)h}^{nh} Y_u du$  by some numerical integration scheme needing only the values of the process on this finer grid. Since the approximations of  $\Delta L^{(h)}$  become thus closer and closer to the true increment as the numerical integration scheme becomes more exact, in [14] the authors derive their asymptotic results when both the observation interval and the observation frequency go to infinity. Note that in practice one does not know a, so one has to estimate it first, which could, e.g., be done by the already described quasi-maximum likelihood approach.

Turning to an example, let us consider the OU process given by

$$dX_t = -0.6X_t dt + dL_t, \tag{6.4}$$

with L being a standardized Gamma process, i.e., Lt having density

$$f_{L_t}(x) = \frac{\gamma^{1/2\gamma t}}{\Gamma(\gamma t)} x^{\gamma t-1} e^{-x\gamma^{1/2}} \mathbf{1}_{[0,\infty)},$$

and the parameter  $\gamma$  being set to 2.

In [31], 100 paths of this OU process on the time interval [0, 5000] have been simulated and then the Lévy increments over time intervals of unit length have been approximated by sampling the OU process over a grid of size h.

In Figure 6 the histogram of the Lévy increments distribution from one path with h = 0.01 is shown, together with the true probability density of L<sub>1</sub>.

If one further averages over all 100 paths, which is equivalent to looking at one path over a time horizon 100 times longer, the fit of the histogram to the true density becomes visually almost perfect (see Figure 7).

Based on the approximate Lévy increments one can now estimate the parameter  $\gamma$  by maximum likelihood. Table 2 shows summary statistics of the resulting estimator for different sampling grid sizes h. The data in the table are based on estimating  $\gamma$  separately for each of the 100 simulated paths.

To conclude, the simulation study illustrates that both the recovery of the background driving Lévy process and the parametric estimation based on approximate increments work quite well.

Foreword

**Featured Essay** 

**Author Profile** 

**Primary Sources** 

About





Figure 6. Probability density of the increments of the standardized Lévy process with  $\gamma = 2$  and the histogram of the estimated increments from one path of the OU process, obtained by sampling the process with grid length 0.01. (Source: [31].)

Foreword

**Featured Essay** 

**Author Profile** 

About Primary Sources



Robert Stelzer: CARMA processes driven by non-Gaussian noise

Figure 7. Probability density of the increments of the standardized Lévy process with  $\gamma = 2$  and the histogram of the estimated increments for all 100 paths of the OU process, obtained by sampling the process with grid length 0.01. (Source: [31].)

#### 7 Concluding Remarks

Finally, we would like to mention that there are other stochastic models, such as the so-called ECOGARCH process of [33] and [34], where both CARMA processes and extensions of CARMA processes are important ingredients. One type of extension is fractionally integrated CARMA (FICARMA) processes (see [13] and [41]). While CARMA processes have an exponentially decaying autocovariance function and thus always have short memory, FICARMA processes exhibit polynomially decaying autocovariance functions and are thus able to model long-memory phenomena (see [21] or [52] for detailed introductions into the topic of long-range dependence). However, the paths of FICARMA processes are continuous. The supOU processes (see [4, 5] and [26]) represent a class of processes that can exhibit long memory and jumps in the paths and are

Featured Essay

Foreword

**Author Profile** 

About Primary Sources Robert Stelzer: CARMA processes driven by non-Gaussian noise

Table 2. Estimated parameters of the standardised driving Lévy process based on 100 paths on  $\left[0,5000\right]$  of the Gamma-driven OU process.

h	Parameter	Sample mean of estimator	Sample standard deviation of estimator
0.01	γ	2.0039	0.0314
0.1	γ	2.0043	0.0340
1	γ	1.9967	0.0539

related to CARMA processes. As noted in [5], multivariate supOU processes can be straightforwardly extended to obtain so-called supCARMA processes. Long memory is (believed to be) encountered in data from many different areas, e.g., finance and telecommunications. Since it is an asymptotic property, and similar effects in the autocorrelation function might be caused by structural breaks (non-stationarity), it is often hardly debated whether there truly is long memory in a time series. The first scientific study considering long-range dependence properties was looking at the water level of the river Nile (see [57]).

From the overview on CARMA processes presented in this paper, it should not only be clear that they are useful in many applications, but also that there are still many questions to be addressed in future research. These include alternatives to the estimators presented here, estimators that work in the heavy-tailed case when one does not have a finite variance or order selection, i.e., a theory on how to choose the orders (p, q) of the autoregressive and moving average polynomial when one fits CARMA processes to observed time series.

#### **Acknowledgements**

The author takes pleasure in thanking Florian Fuchs, Claudia Klüppelberg and Eckhard Schlemm for comments on previous drafts and Maria Graf for allowing him to use material from her diploma thesis in Section 6.2. Financial support from the TUM Institute for Advanced Study funded by the German Excellence Initiative through a Carl-von-Linde Junior Fellowship is gratefully acknowledged.

**Featured Essay** 

**Author Profile** 

About Primary Sources Robert Stelzer: CARMA processes driven by non-Gaussian noise

#### **Bibliography**

- D. Applebaum. Lévy Processes and Stochastic Calculus, volume 93 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 2004.
- [2] M. Arató. Linear Stochastic Systems with Constant Coefficients, volume 45 of Lecture Notes in Control and Information Sciences. Springer, Berlin, 1982.
- [3] O. E. Barndorff-Nielsen. Processes of normal inverse Gaussian type. *Finance Stoch.*, 2:41–68, 1998.
- [4] O. E. Barndorff-Nielsen. Superposition of Ornstein–Uhlenbeck type processes. *Theory Probab. Appl.*, 45:175–194, 2001.
- [5] O. E. Barndorff-Nielsen and R. Stelzer. Multivariate supOU processes. *Ann. Appl. Probab.*, 21:140–182, 2011.
- [6] J. Bertoin. *Lévy Processes*. Cambridge University Press, Cambridge, 1998.
- [7] P. Blæsild and J. L. Jensen. Multivariate distributions of hyperbolic type. In C. Taillie, G. P. Patil, and B. A. Baldessari, editors, *Statistical* distributions in scientific work - Proceedings of the NATO Advanced Study Institute held at the Université degli Studi di Trieste, Triest, Italy, July 10 - August 1, 1980, volume 4, pages 45–66, Dordrecht, Holland, 1981. D. Reidel.
- [8] P. J. Brockwell. Lévy-driven CARMA processes. Ann. Inst. Statist. Math., 52:1–18, 2001.
- [9] P. J. Brockwell. Continuous-time ARMA processes. In D. N. Shanbhag and C. R. Rao, editors, *Stochastic Processes: Theory and Methods*, volume 19 of *Handbook of Statistics*, pages 249–276, Amsterdam, 2001. Elsevier.
- [10] P. J. Brockwell. Lévy-driven continuous-time ARMA processes. In T. G. Andersen, R. Davis, J.-P. Kreiß, and T. Mikosch, editors, *Handbook of Financial Time Series*, pages 457 – 480, Berlin, 2009. Springer.
- [11] P. J. Brockwell and R. A. Davis. *Time Series: Theory and Methods*. Springer, New York, 2nd edition, 1991.

**Featured Essay** 

**Author Profile** 

About Primary Sources Robert Stelzer: CARMA processes driven by non-Gaussian noise

[12] P. J. Brockwell and A. Lindner. Existence and uniqueness of stationary Lévy-driven CARMA processes. *Stoch. Process. Appl.*, 119:2660–2681, 2009.

- [13] P. J. Brockwell and T. Marquardt. Lévy-driven and fractionally integrated ARMA processes with continuous time parameter. *Statist. Sinica*, 15:477–494, 2005.
- [14] P. J. Brockwell and E. Schlemm. Parametric estimation of the driving Lévy process of multivariate CARMA processes from discrete observations. submitted for publication, 2011.
- [15] P. J. Brockwell, R. A. Davis, and Y. Yang. Continuous-time Gaussian autoregression. *Statist. Sinica*, 17:63–80, 2007.
- [16] P. J. Brockwell, R. A. Davis, and Y. Yang. Estimation for nonnegative Lévy-driven Ornstein-Uhlenbeck processes. J. Appl. Probab., 44: 977–989, 2007.
- [17] P. J. Brockwell, R. A. Davis, and Y. Yang. Estimation of non-negative Lévy-driven CARMA processes. J. Bus. Econom. Statist., to appear, 2011.
- [18] P. J. Brockwell, V. Ferrazzano, and C. Klüppelberg. High frequency sampling of a continuous-time ARMA process. 2011, Journal of Time Series Analysis, to appear.
- [19] J. L. Doob. The elementary Gaussian processes. Ann. Math. Stat., 25: 229–282, 1944.
- [20] A. Doucet, N. de Freitas, and N. J. Gordon, editors. Sequential Monte Carlo Methods in Practice, Statistics for Engineering and Information Science, New York, 2001, Springer.
- [21] P. Doukhan, M. S. Taqqu, and G. Oppenheim, editors. *Long-Range Dependence*, Boston, 2003. Birkhäuser.
- [22] P. Embrechts, C. Klüppelberg, and T. Mikosch. *Modelling Extremal Events for Insurance and Finance*. Springer, Berlin, 1997.
- [23] V. Fasen. Extremes of regularly varying Lévy-driven mixed moving average processes. Adv. in Appl. Probab., 37:993–1014, 2005.
- [24] V. Fasen. Extremes of subexponential Lévy driven moving average processes. *Stoch. Proc. Appl.*, 116:1066–1087, 2006.

**Featured Essay** 

**Author Profile** 

About Primary Sources Robert Stelzer: CARMA processes driven by non-Gaussian noise

[25] V. Fasen. Extremes of Lévy driven mixed MA processes with convolution equivalent distributions. *Extremes*, 12:265–296, 2009.

- [26] V. Fasen and C. Klüppelberg. Extremes of supOU processes. In F. E. Benth, G. Di Nunno, T. Lindstrom, B. Øksendal, and T. Zhang, editors, *Stochastic Analysis and Applications: The Abel Symposium* 2005, volume 2 of *Abel Symposia*, pages 340–359, Berlin, 2007. Springer.
- [27] F. Fuchs and R. Stelzer. Spectral representation of multivariate regularly varying Lévy and CARMA processes. J. Theoret. Probab., to appear, 2011.
- [28] F. Fuchs and R. Stelzer. Mixing conditions for multivariate infinitely divisible processes with an application to mixed moving averages and the supOU stochastic volatility model. ESAIM Probability Statist., to appear, 2011.
- [29] I. García, C. Klüppelberg, and G. Müller. Estimation of stable CARMA models with an application to electricity spot prices. *Stat. Model.*, 11: 447–470, 2011.
- [30] H. Garnier and L. Wang, editors. *Identification of Continuous-time Models from Sampled Data*. Advances in Industrial Control. Springer, London, 2008.
- [31] M. Graf. Parametric and nonparametric estimation of positive Ornstein-Uhlenbeck type processes. Diploma thesis, Centre for Mathematical Sciences, TU München, 2009. Available from http://www-m4.ma.tum.de/Diplarb.
- [32] E. J. Hannan and M. Deistler. *The statistical theory of linear systems*. Wiley Series in Probability and Mathematical Statistics: Probability and Mathematical Statistics. John Wiley & Sons, New York, 1988. ISBN 0-471-80777-X.
- [33] S. Haug and C. Czado. An exponential continuous time GARCH process. J. Appl. Probab., 44: 960–976, 2007.
- [34] S. Haug and R. Stelzer. Multivariate ECOGARCH processes. *Econometric Theory*, 27: 344-371, 2011.
- [35] B. Jørgensen. *Statistical Properties of the Generalized Inverse Gaussian Distribution*. Lecture Notes in Statistics. Springer, New York, 1982.

**Featured Essay** 

**Author Profile** 

About Primary Sources Robert Stelzer: CARMA processes driven by non-Gaussian noise

[36] Z. J. Jurek and D. J. Mason. Operator-limit Distributions in Probability Theory. John Wiley & Sons, New York, 1993.

- [37] A. E. Kyprianou. Introductory Lectures on Fluctuations of Lévy Processes with Applications. Universitext. Springer, Berlin, 2006.
- [38] E. K. Larsson. Limiting sampling results for continuous-time ARMA systems. *Internat. J. Control*, 78:461–473, 2005.
- [39] E. K. Larsson, M. Mossberg, and T. Söderström. An overview of important practical aspects of continuous-time ARMA system identification. *Circuits Systems Signal Process.*, 25:17–46, 2006.
- [40] H. Lütkepohl. *New Introduction to Multiple Time Series Analysis*. Springer, Berlin, 2005.
- [41] T. Marquardt. Multivariate fractionally integrated CARMA processes. *J. Multivariate Anal.*, 98:1705–1725, 2007.
- [42] T. Marquardt and R. Stelzer. Multivariate CARMA processes. *Stochastic Process. Appl.*, 117:96–120, 2007.
- [43] T. Mikosch, S. I. Resnick, R. Holger, and A. Stegeman. Is network traffic approximated by stable Lévy motion or fractional Brownian motion? *Ann. Appl. Probab.*, 12:23–68, 2002.
- [44] M. Moser and R. Stelzer. Tail Behavior of Multivariate Lévy-Driven Mixed Moving Average Processes and supOU Stochastic Volatility Models. Adv. Appl. Probab., to appear, 2011.
- [45] M. Mossberg and E. K. Larsson. Fast and approximative estimation of continuous-time stochastic signals from discrete-time data. In *IEEE International Conference on Acoustics, Speech, and Signal Processing, 2004. Proceedings. (ICASSP '04)*, volume 2, pages 529–532, Montreal, QC, May 17-21, 2004, 2004.
- [46] K. Prause. The Generalized Hyperbolic Model: Estimation, Financial Derivatives and Risk Measures. Dissertation, Mathematische Fakultät, Albert-Ludwigs-Universität Freiburg i. Br., Freiburg, Germany, 1999.
- [47] M. B. Priestley. *Spectral analysis and time series. Vol. 1 & 2.* Academic Press, London, 1981.
- [48] P. Protter. Stochastic Integration and Differential Equations, volume 21 of Stochastic Modelling and Applied Probability. Springer-Verlag, New York, 2nd edition, 2004.

**Featured Essay** 

**Author Profile** 

About Primary Sources Robert Stelzer: CARMA processes driven by non-Gaussian noise

- [49] S. I. Resnick. *Extreme Values, Regular Variation and Point Processes*. Springer, New York, 1987.
- [50] S. I. Resnick. Heavy tail modeling and teletraffic data: special invited paper. *Ann. Statist.*, 25:1805–1869, 1997.
- [51] S. I. Resnick. Heavy-Tail Phenomena. Springer, 2007.

- [52] G. Samorodnitsky. Long range dependence. *Found. Trends Stoch. Syst.*, 1:163–257, 2006.
- [53] K. Sato. Lévy Processes and Infinitely Divisible Distributions, volume 68 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 1999.
- [54] K. Sato and M. Yamazato. Operator-selfdecomposable distributions as limit distributions of processes of Ornstein-Uhlenbeck type. *Stochastic Process. Appl.*, 17:73–100, 1984.
- [55] E. Schlemm and R. Stelzer. Multivariate CARMA processes, continous-time state space models and complete regularity of the innovations of the sampled processes. *Bernoulli*, 2011. accepted for publication.
- [56] E. Schlemm and R. Stelzer. Quasi maximum likelihood estimation for strongly mixing state space models and multivariate CARMA processes, submitted for publication, 2011.
- [57] O. Tousson. Mémoire sur l'Histoire du Nil, Mémoires de l'Institut d'Egypte (Cairo), 1925.

TUM-IAS Primary Sources Author Profile

# Author profile: Robert Josef Stelzer

7

e halls bein horsing

d. Ordnung

#### **Featured Essay**

Author Profile

About Primary Sources

The paper at the heart of our inaugural issue springs from a prolific source.

The impressive number of papers and talks generated by Robert Stelzer's research during the past few years, in what must still be called his "early career," is just one measure of his scientific creativity. Already with his doctoral thesis, "Multivariate Continuous Time Stochastic Volatility Models Driven by a Lévy Process," Stelzer won recognition for the quality and originality of his work.

In 2008 he received the prestigious prize for outstanding dissertations awarded once every two years by the Probability and Statistics Group of the Deutsche Mathematiker Vereinigung.

Add to that the broad set of topics he has addressed, a deep level of engagement with students and collaborators, and his extensive professional activities: Stelzer has organized both tightly focused and broadly interdisciplinary scientific meetings, and he has been editor of the newsletter of the Bernoulli Society for the last two years. He is a member of the Applied Probability Society, the **Institute of Mathematical** Statistics, the DMV, and its Probability and Statistics Group. He serves as associate editor for two journals and as managing editor for a small book series, as a referee for papers submitted to many journals, and as an expert reviewer for the Alexander von Humboldt Foundation.

A postdoc at the Technische Universität München when his Carl von Linde Junior Fellowship at the TUM-IAS began, Robert Stelzer recently accepted a call — at the age of 30 — to become a professor at Ulm University, and the director of its Institute of Mathematical Finance.

His research interests include financial mathematics, multivariate stochastic volatility models, stochastic processes, Lévy processes, multivariate time series analysis, random matrices, Markov switching models, and extreme value theory. One characteristic these diverse fields have in common is that advances in theory and methodology can suggest or directly provide useful tools for real-world applications. And that, according to Stelzer, goes a long way toward explaining the greatest challenges he and his colleagues face, as well as their most compelling motivations.

#### TUM-IAS Primary Sources Author Profile

Featured Essay

### Author Profile

About Primary Sources



#### TUM-IAS Primary Sources Author Profile

**Featured Essay** 

#### Author Profile

About Primary Sources



"For us," he says, "there definitely is a kind of beauty in the mathematical theory itself, in the expressive power that enables researchers to pose and explore such a wide range of problems. It can sometimes be hard to develop a common language for collaboration with experts from other fields; but once you have that basis for

communication, the theoretical work can have a strong practical impact, which is a great motivation for the theoretical work. Likewise, 'simple' questions arising in applications may well go beyond what is understood so far and thus lead to challenging new research questions."

#### **Photo Credits:**

Page 1, © Astrid Eckert / TU München Page 3, © Reinhard Sandbothe / Pixelio Page 5, Fractal Image from Wiki Commons Page 42, © Andreas Heddergott / TU München Page 44, 45, © Faces by Frank / TU München Page 46, Fractal Images from Wiki Commons Photos © Astrid Eckert, Andreas Heddergott / TU München

#### TUM-IAS Primary Sources Author Profile













### **About Primary Sources**

Primary Sources is a research journal published by the TUM Institute for Advanced Study to stimulate interdisciplinary communication and innovation at the highest levels of science and engineering.

The TUM-IAS, a central institute of the Technische Universität München, was created within the framework of the Excellence Initiative of Germany's federal and state governments. Technische Universität München is one of Germany's leading universities, with a research and teaching portfolio spanning the engineering sciences, natural sciences, life sciences, medicine, and economic sciences.

Copyright © 2011 Technische Universität München Institute for Advanced Study Lichtenbergstraße 2 a 85748 Garching, Germany

• For more information, please visit and bookmark our home pages: www.tum.de and www.tum-ias.de.

• To propose or submit an essay, please contact Prof. Patrick Dewilde (dewilde@zv.tum.de), or Patrick Regan (regan@zv.tum.de).



Patrick Dewilde TUM-IAS Director, Scientific Editor



Patrick Regan Editor



Stefanie Hofmann Managing Editor



Tatjana Steinberger Managing Editor

Design & LaTeX Support: Amy Hendrickson, TeXnology Inc.



