

Roman Andreev & Christine Tobler. “Multilevel preconditioning and low-rank tensor iteration for space-time simultaneous discretizations of parabolic PDEs”. In: *Numer. Linear Algebra Appl.* 22.2 (2015), pp. 317–337. ISSN: 1070-5325. DOI: 10.1002/nla.1951. URL: <http://dx.doi.org/10.1002/nla.1951>.

Tammo Jan Dijkema & Rob Stevenson. “A sparse Laplacian in tensor product wavelet coordinates”. In: *Numer. Math.* 115.3 (2010), pp. 433–449. ISSN: 0029-599X. DOI: 10.1007/s00211-010-0288-5. URL: <http://dx.doi.org/10.1007/s00211-010-0288-5>.

Ballani & Grasedyck. “A projection method to solve linear systems in tensor format”. In: *Numer. Linear Algebra Appl.* 20.1 (2013), pp. 27–43. ISSN: 1070-5325. DOI: 10.1002/nla.1818. URL: <http://dx.doi.org/10.1002/nla.1818>.

Billaud-Friess, Nouy, & Zahm. “A tensor approximation method based on ideal minimal residual formulations for the solution of high-dimensional problems”. In: *ESAIM: Mathematical Modelling and Numerical Analysis* 48 (06 Nov. 2014), pp. 1777–1806. ISSN: 1290-3841. DOI: 10.1051/m2an/2014019. URL: http://www.esaim-m2an.org/article_S0764583X14000193.

Hackbusch et al. “Use of tensor formats in elliptic eigenvalue problems”. In: *Numer. Linear Algebra Appl.* 19.1 (2012), pp. 133–151. ISSN: 1070-5325. DOI: 10.1002/nla.793. URL: <http://dx.doi.org/10.1002/nla.793>.

Khoromskij. “Tensor-structured preconditioners and approximate inverse of elliptic operators in \mathbb{R}^d ”. In: *Constr. Approx.* 30.3 (2009), pp. 599–620. ISSN: 0176-4276. DOI: 10.1007/s00365-009-9068-9. URL: <http://dx.doi.org/10.1007/s00365-009-9068-9>.