### ARTICLE

Chronic lymphocytic leukemia



# Short telomeres are associated with inferior outcome, genomic complexity, and clonal evolution in chronic lymphocytic leukemia

Billy Michael Chelliah Jebaraj<sup>1</sup> · Eugen Tausch<sup>1</sup> · Dan A. Landau<sup>2,3,4</sup> · Jasmin Bahlo<sup>5</sup> · Sandra Robrecht<sup>5</sup> · Amaro N. Taylor-Weiner<sup>6</sup> · Johannes Bloehdorn<sup>1</sup> · Annika Scheffold<sup>1</sup> · Daniel Mertens<sup>1,7</sup> · Sebastian Böttcher<sup>8,12</sup> · Michael Kneba<sup>8</sup> · Ulrich Jäger<sup>9</sup> · Thorsten Zenz<sup>1,13</sup> · Michael K. Wenger<sup>10</sup> · Guenter Fingerle-Rowson<sup>10</sup> · Clemens Wendtner<sup>11</sup> · Anna-Maria Fink<sup>5</sup> · Catherine J. Wu<sup>6</sup> · Barbara Eichhorst<sup>5</sup> · Kirsten Fischer<sup>5</sup> · Michael Hallek<sup>5</sup> · Hartmut Döhner<sup>1</sup> · Stephan Stilgenbauer<sup>1</sup>

Received: 23 September 2018 / Revised: 19 January 2019 / Accepted: 11 February 2019 © Springer Nature Limited 2019

### Abstract

Telomere length in chronic lymphocytic leukemia (CLL) has been shown to be of prognostic importance, but the analyses have largely been executed on heterogeneous patient cohorts outside of clinical trials. In the present study, we performed a comprehensive analysis of telomere length associations in the well characterized CLL8 trial (n = 620) of the German CLL study group, with validation in a representative cohort of the CLL4 trial (n = 293). Absolute telomere length was analyzed using quantitative-PCR. Apart from identifying associations of short telomere length with adverse prognostic factors and survival, the study identified cases with 17p- and 11q- associated with *TP53* and *ATM* loss, respectively, to have the shortest telomeres, even when these aberrations were present in small subclones. Thus, telomere shortening may precede acquisition of the high-risk aberrations, contributing to disease evolution. In line with this, telomere shortening was associated with an increase in genomic complexity as well as clonal evolution, highlighting its importance as a biomarker especially in monitoring disease progression in non-high-risk CLL.

**Supplementary information** The online version of this article (https://doi.org/10.1038/s41375-019-0446-4) contains supplementary material, which is available to authorized users.

Stephan Stilgenbauer stephan.stilgenbauer@uniklinik-ulm.de

- <sup>1</sup> Department of Internal Medicine III, Ulm University, Ulm, Germany
- <sup>2</sup> Department of Medicine, Weill Cornell Medicine, New York, NY, USA
- <sup>3</sup> Department of Physiology and Biophysics, Weill Cornell Medicine, New York, USA
- <sup>4</sup> New York Genome Center, New York, NY, USA
- <sup>5</sup> Department I for Internal Medicine and Centre for Integrated Oncology, University of Cologne, Cologne, Germany
- <sup>6</sup> Broad Institute, Cambridge, MA, USA

# Introduction

Chronic lymphocytic leukemia (CLL) has a variable clinical course ranging from indolent to high-risk disease with short survival despite intensive treatment. Prognostication, therefore, is of key importance, and various factors such as clinical characteristics, immunoglobulin heavy chain gene (IGHV) mutation status, genomic aberrations, and gene

- <sup>7</sup> Mechanisms of Leukemogenesis, DKFZ, Heidelberg, Germany
- <sup>8</sup> Department II of Internal Medicine, University Hospital of Schleswig-Holstein, Kiel, Germany
- <sup>9</sup> Department of Medicine I, Division of Hematology and Hemostaeology, Medical University of Vienna, Vienna, Austria
- <sup>10</sup> F. Hoffmann-La Roche Ltd, Basel, Switzerland
- <sup>11</sup> Klinikum Schwabing, Academic Teaching Hospital of University of Munich, Munich, Germany
- <sup>12</sup> Present address: Division of Internal Medicine, Medical Clinic III, Rostock University Medical Center, Rostock, Germany
- <sup>13</sup> Present address: Division of Hematology, University Hospital Zürich, Zürich, Switzerland

mutions have been proposed [1, 2]. Also, telomere length in CLL has been associated with biological and clinical disease characteristics as well as outcome [3-9].

Telomeres are nucleoprotein complexes at the ends of each chromosome whose main function is capping of chromosomal ends to prevent them from being recognized as DNA damage. In somatic cells, telomeres shorten with cell division owing to the end-replication problem. Cells with short telomere length undergo senescence and upon abrogation of the senescence checkpoints, the cells can undergo proliferation with further telomere shortening until the crisis stage leading to cell death. In stem cells and germ cells, telomere length is maintained by the enzyme complex telomerase [10–12].

Telomere dysfunction is described to be associated with tumorigenesis across various cancers [13–19] and short telomeres may favor genomic instability, checkpoint deregulation, and activation of telomerase, leading to malignant transformation [12]. Different treatment strategies exploit the crucial role of telomere dysfunction, and targeting telomerase dependence of cancers shows promising impact in clinical trials [20]. In CLL, cases with short telomeres are associated with high telomerase expression and activity, presumably to maintain the critical telomere length for the survival of the cell. Telomerase expression and activity along with telomere length were shown to be of prognostic value in CLL [3, 4, 8]. However in CLL, the impact of drugs targeting telomerase is more difficult to assay, as the cells show limited proliferation in vitro.

The prognostic value of telomere length in CLL has mainly been tested on small and heterogeneous patient cohorts. Here, we performed a comprehensive analysis of telomere length associations with disease characteristics and outcome using the large, well-characterized CLL8 trial of the German CLL study group (GCLLSG) with mature follow-up [21]. The findings were validated using the CLL4 trial of the GCLLSG [22].

The associations of short telomere length with poor prognostic factors in CLL are largely presumed to be a consequence of the underlying genetic characteristics. Here, we verified if telomere shortening, in addition to being a marker of cell proliferation, could have a role in acquisition of high-risk aberrations, favoring genomic complexity and clonal evolution.

# Materials and methods

### Patient cohort

The CLL8 trial was an international prospective, multicenter, randomized phase III trial for first-line treatment comparing fludarabine and cyclophosphamide (FC) with FC and rituximab (FCR) as previously reported [23]. In total, 817 patients in need for treatment were enrolled. Among

these, 620 (76%) cases with available DNA were included in the analysis of telomere length. In terms of validation, the CLL4 trial of the GCLLSG was used, which included 362 patients that were randomized to first-line treatment with fludarabine alone (F) vs. FC [22]. In all, 293 (81%) cases with available DNA were included for telomere-length analysis. The analyzed cohorts were representative of the whole trial populations (Suppl. Table 1). CLL4 and CLL8 were open-label studies without blinding (investigators and study personnel were not blinded to treatment assessment). B cells from age-matched healthy probands (n = 20) were isolated from peripheral blood samples by CD19 sorting and used as control. The study was conducted in accordance with the Declaration of Helsinki, approved by our institutional review board, and informed consent was obtained from all patients before inclusion in the study.

#### **Telomere length measurement using Q-PCR**

Telomere length measurement was carried out using a quantitative-PCR (Q-PCR)-based technique, favouring measurement of large numbers of samples with relatively low quantities of DNA [24]. The primers used to amplify telomere and single-copy genes (SCG) were tel1b, tel2b, and HBG3, HBG4, respectively [24]. The absolute telomere length was obtained by using synthetic oligonucleotide standards for telomere (84 bp) and SCG (81 bp) PCR. In brief, a serial dilution of the telomere and SCG standard was prepared and the amount of DNA molecules in each standard was calculated as previously described [25]. In total, 12 ng of DNA was used per reaction (total volume of 10 µl) in triplicates for the telomere and SCG PCRs and amplified using Qiagen quantitect SYBR green (Qiagen, Hilden, Germany) in 384-well plates and analyzed using 7900HT fast real-time PCR system (Applied Biosystems, Darmstadt, Germany). The Q-PCR technique was validated by terminal restriction fragment length (TRF) analysis. Six telomere length controls with known telomere length, analyzed using TRF analysis were included in every plate to detect variation. The TRF value of telomere length for each sample was calculated from the linear regression of Q-PCR versus TRF.

#### Validation of Q-PCR technique with TRF analysis

The Q-PCR technique was validated by TRF analysis and Southern hybridization. A total of  $6 \mu g$  of non-degraded DNA was digested overnight using *Hin*fI and *Rsa*I (New England Biolabs, Frankfurt, Germany) and resolved on a 0.8% agarose gel. In gel hybridization was carried out by drying the gel and hybridizing with a telomere-specific probe, end labeled with <sup>32</sup>P. The mean telomere length was analyzed from the autoradiograph as previously described [26]. A correlation of  $R^2 = 0.8516$  was obtained upon comparison of telomere length measured using Q-PCR and TRF in a control sample set (n = 18) (Suppl. Fig. 1a).

#### **Genetic analyses**

Genomic aberrations were analyzed using fluorescence in situ hybridization (FISH) [27] and IGHV status [28] by Sanger sequencing, respectively. Mutations in TP53 were identified using Amplichip (Roche Molecular Systems, Mannheim, Germany) and confirmed by Sanger sequencing (Big Dye Terminator Kit and ABI 3100 sequencer; Applied Biosystems, Darmstadt, Germany). NOTCH1 mutations were identified using Sanger sequencing and SF3B1 mutations were identified using dHPLC on a WAVE 3500HT System (Transgenomic Inc., Crewe, UK) and confirmed by Sanger sequencing [29]. Detection of copy number alteration and of loss of heterozygosity was performed by singlenucleotide polymorphism (SNP) array analysis (Affymetrix 6.0., Santa Clara, CA, USA) [30]. Whole-exome sequencing (WES) was performed on HiSeq 2000 (Illumina, San Diego, CA, USA), as previously described [31].

#### **Statistical analyses**

Associations between patient characteristics and telomere length were evaluated both as grouped variable (with median as cutoff) and as a continuous variable using chi-square test and non-parametric rank-sum test as appropriate. Correlations of telomere length with genomic complexity, CLL drivers and clone size of aberrations were analyzed using Spearman rank correlation  $(r_s)$  or linear regression  $(R^2)$ , as appropriate. Timeto-event parameters included progression free survival (PFS) and overall survival (OS). Kaplan-Meier estimates and twosided non-stratified log-rank tests were used for analyzing survival times. Hazard ratios (HR) and confidence intervals (CI) were calculated using Cox proportional-hazard regression analyses. Multivariable analyses were performed using Cox regression with stepwise selection including only those variables that were significantly associated with PFS or OS in univariate analyses. All statistical tests were two-sided and a P value of < 0.05 was considered as significant. No adjustments for multiple testing were done. The analyses were performed using SPSS 23.0 (SPSS, Chicago, IL) and Graphpad prism (versions 5.0 and 8.0.2).

## Results

# Telomere length variation in CLL is largely restricted to tumor cells

To investigate whether telomere shortening in CLL is restricted to the tumor cell fraction, 48 cases from CLL8

with paired tumor (CD19 +), non-tumor (CD19 -), and unsorted samples were analyzed. Also, CD19 + samples were included from 20 age-matched healthy donors (median: 56 years), out of which eight cases had a paired CD19- sample. The median telomere length of CLL cells (CD19+) was significantly shorter compared with the non B cells (CD19-) of CLL patients (median: 3.37 kb vs. 5.42 kb; P = 0.0004), whereas, in healthy donors, nonmalignant B cells had significantly longer telomeres than their CD19- counterparts in each case (7.54 kb vs. 5.40 kb, respectively; P < 0.001) (Fig. 1a). Telomere length of tumor B cells significantly differed from that of normal B cells (median: 3.37 kb vs. 7.54 kb, P < 0.0001), whereas the CD19- cells from CLL patients and healthy donors had similar telomere length distribution (5.42 kb vs. 5.40 kb). Further, the influence of the disease features on telomere length was apparent in the CD19 + tumor cell fraction where a correlation with patient's age was absent ( $R^2 = 0.002$ , P = 0.919) (Suppl. Fig. 1b). In the non-tumor fraction, an association of telomere length with age was retained  $(R^2 = 0.191, P = 0.002)$  (Suppl. Fig. 1c).

CD19 enriched samples were available only for a subset of cases hence in the analysis of the full CLL8 and CLL4 cohorts, DNA from either the CD19 sorted or the unsorted cells were used for telomere length measurement. Of note, telomere length was similar between the CD19 + sorted and unsorted samples from same patients (median: 3.38 kb vs. 3.48 kb; P = 0.197) (Fig. 1a). Absence of adverse influence of non-tumor fraction on telomere length associations in samples with low WBC count (<  $50 \times 10^9$  cells/L) without CD19 enrichment (N = 128) was verified by analyzing correlation of telomere length with WBC count as well as survival (Suppl. Fig. 2).

### Telomere length in CLL8 and its association with clinical characteristics

The telomere length of the CLL8 samples (n = 620) was found to be highly variable, ranging from 1.94 kb to 33.56 kb (median 4.56 kb). Analysis of the frequency distribution of the telomere lengths revealed a skewing toward shorter telomeres, with 46.9% of the cases represented in the telomere length range of  $\ge 3$  kb and < 5 kb. In total, 11.7% of the cases analyzed had very short telomeres of < 3 kb. A subset of CLL cases (1.6%) displayed extremely long telomeres of  $\ge 15$  kb (15.57–33.56 kb) (Suppl. Fig. 3a) and the telomere lengths obtained by q-PCR were validated using TRF for cases with available DNA with a strong correlation ( $R^2 = 0.981$ , P < 0.001) (Suppl. Fig. 3b, c). Although an overestimation of telomere length by q-PCR was observed in increasing telomere lengths, these cases had significantly



**Fig. 1 a** Comparison of telomere length between CD19+, CD19-, and unsorted CLL samples (n = 48) and with age-matched healthy donors n = 20 (CD19+) and n = 8 (CD19-). **b** telomere length in cases with mutated and unmutated IGHV; mutated and wildtype *TP53*, *NOTCH1*, *SF3B1*. **c** Telomere length in the hierarchically classified genomic aberration subgroups namely, 17p-, 11q-, + 12q, normal karyotype and 13q- in CLL8 (red bars indicate the median telomere length). All *P* values from Mann–Whitney test

longer telomeres compared with that of normal B cells as measured by TRF (P < 0.001).

Associations of telomere length and clinical characteristics in the CLL8 cohort were analyzed by comparing groups defined by short and long telomeres (cutoff at the median of 4.56 kb). Telomere length showed no significant association with age (P = 0.629), sex (P = 0.168), presence of B-symptoms (P = 0.297), Eastern Cooperative Oncology Group (**ECOG**) status (P = 0.224), and high serum beta-2 microglobulin (s- $\beta_2$ MG > 3.5; P = 0.653). In contrast, short telomeres were significantly associated with other adverse risk characteristics such as high WBC ( $\geq$  50 Giga/L; P < 0.001), high serum thymidine kinase (s-TK > 10.0 U/L; P < 0.001) and Binet stage B (P = 0.002) (Table 1), whereas longer telomeres were associated with Binet C stage (P = 0.007).

# Association of telomere length with genetic characteristics

Among the genetic characteristics, unmutated IGHV, mutated TP53, and mutated SF3B1 were significantly associated (P <0.001) with short telomeres, whereas NOTCH1 (P = 0.696) mutations were not (Suppl. Table 2). The hierarchically categorized genomic aberration subgroups 17p- and 11q- had significantly shorter telomeres (both P < 0.001), whereas cases with 13q- as the sole abnormality (P < 0.001) and no abnormality (P = 0.020) had longer telomeres. The presence of + 12q (P = 0.071) showed no association with telomere length (Suppl. Table 2). Figure 1b, c show the distribution of telomere length (as continuous variable) in the different gene mutation and genomic aberration subgroups. ZAP70 + (>20%) cases were found to be significantly associated with short telomeres (P = 0.010). All cases with very long telomeres belonged to the favorable prognostic subgroup, mutated IGHV, predominantly with 13q- as sole abnormality (70%).

# Impact of telomere length on clinical response, MRD, and survival

Though significantly associated with adverse clinical and genetic characteristics, telomere length in CLL8 had no association with clinical response (CR: 37.7% vs. 41.2% and PR: 53.2% vs. 52.4%; P = 0.400) (Suppl. Table 3). Also, telomere length did not impact MRD response in peripheral blood (P = 0.751) and bone marrow (P = 0.213) (Suppl. Table 4). At a median observation time of 69.7 months, there were 422 (FC 228 vs. FCR 194) events for PFS and 208 (FC 117 vs. FCR 91) for OS. The short telomere subgroups ( $\leq$  median) had significantly shorter PFS in both the FC and FCR treatment arms (FC: 27.1 vs.

Short telomeres are associated with inferior outcome, genomic complexity, and clonal evolution in...

 
 Table 1 Correlation of telomere length (TL) with clinical characteristics (*P* value from chi-souare test)

| Patient characteristics        | CLL8              |                  |           | CLL4       |                  |         |  |
|--------------------------------|-------------------|------------------|-----------|------------|------------------|---------|--|
|                                | Short TL<br>N (%) | Long TL<br>N (%) | long TL P |            | Long TL<br>N (%) | Р       |  |
| Analysis cohort, N             | 310               | 310              |           | 146        | 147              |         |  |
| Age at entry (years)           | 310               | 310              |           | 146        | 147              |         |  |
| >60                            | 169 (54.5)        | 163 (52.6)       | 0.629     | 56 (38.4)  | 44 (29.9)        | 0.128   |  |
| Sex                            | 310               | 310              |           | 146        | 147              |         |  |
| Male                           | 238 (76.8)        | 223 (71.9)       | 0.168     | 113 (77.4) | 107 (72.8)       | 0.362   |  |
| Binet stage                    | 310               | 310              |           | 145        | 147              |         |  |
| А                              | 14 (4.5)          | 20 (6.5)         | 0.009     | 11 (7.6)   | 15 (10.2)        | 0.073   |  |
| В                              | 215 (69.4)        | 178 (57.4)       |           | 71 (49.0)  | 87 (59.2)        |         |  |
| С                              | 81 (26.1)         | 112 (36.1)       |           | 63 (43.4)  | 45 (30.6)        |         |  |
| B-symptoms                     | 310               | 308              |           | 140        | 144              |         |  |
| Yes                            | 134 (43.2)        | 146 (47.4)       | 0.297     | 57 (40.7)  | 62 (43.1)        | 0.689   |  |
| ECOG status                    | 300               | 293              |           | 141        | 137              |         |  |
| ≥1                             | 146 (48.7)        | 128 (43.7)       | 0.224     | 65 (46.1)  | 61 (44.5)        | 0.792   |  |
| WBC count (10 <sup>9</sup> /L) | 304               | 307              |           | 143        | 147              |         |  |
| ≥50                            | 220 (72.4)        | 178 (58.0)       | < 0.001   | 105 (73.4) | 71 (48.3)        | < 0.001 |  |
| s-TK (U/L)                     | 282               | 263              |           | 121        | 131              |         |  |
| >10.0                          | 232 (82.3)        | 177 (67.3)       | < 0.001   | 110 (90.9) | 81 (61.8)        | < 0.001 |  |
| $s-\beta_2 MG \ (mg/L)$        | 282               | 263              |           | 121        | 134              |         |  |
| >3.5                           | 93 (33.0)         | 82 (31.2)        | 0.653     | 62 (51.2)  | 46 (34.3)        | 0.006   |  |

44.0 months, HR 1.934, 95% CI 1.486–2.517, P < 0.001; FCR: 44.8 vs. 69.9 months, HR 1.863, 95% CI 1.395– 2.489, P < 0.001) (Fig. 2a, b). Similarly, short telomeres were significantly associated with shorter OS in both FC and FCR treatments (FC: 70.9 vs. 89.5 months, HR 2.166, 95% CI 1.489–3.151, P < 0.001; FCR: 90.2 months vs. median not reached, HR 1.680, 95% CI 1.098–2.570, P =0.017) (Fig. 2c, d).

Multivariable analysis for PFS was performed including variables that were significant in univariate analyses (treatment arm, gender, genomic aberrations by FISH, mutational status of IGHV, TP53, and SF3B1, s-TK, s-β<sub>2</sub>MG and telomere length categorized based on median). Although FCR treatment (P < 0.001), 17p- (P < 0.001), 11q- (P < 0.001), unmutated IGHV (P < 0.001), mutated TP53 (P = 0.006), SF3B1 mutation (P = 0.018), and s-TK > 10 (P = 0.016)were significant, telomere length was not identified as an independent prognostic factor for PFS (Suppl. Table 5). For OS, treatment arm, age, ECOG status, CIRS score, genomic aberrations by FISH, mutational status of IGHV and TP53, s-TK, s- $\beta_2$ MG, and telomere length categorized based on median were identified as potential candidates in univariate analyses and therefore included as variables for multivariable analysis. Whereas telomere length was not significant, FCR (P = 0.012), ECOG  $\ge 1$  (P = 0.023), CIRS score (P = 0.001), 17p- (P = 0.001), unmutated IGHV (P =0.001), TP53 mutation (P < 0.001), s-TK > 10 (P = 0.020),

and s- $\beta_2$ MG (P = 0.014) were identified as independent prognostic factors (Suppl. Table 6).

# Analysis of associations with telomere length as a continuous variable in CLL8

Associations with different clinical and genetic characteristics were additionally studied considering telomere length as a continuous variable. The results obtained confirmed the observations from categorical (median) analysis (Suppl. Table 7, Fig. 1b, c). Telomere length associations with genomic aberrations within the IGHV subgroups were similar to that of the whole cohort, with 17p- and 11q- associated with the shortest telomere length (Suppl. Fig. 4). When analyzed as a continuous variable, telomere length showed a significant association with PFS (FC: HR 0.921, 95% CI 0.864–0.980 P = 0.010; FCR: HR 0.838, 95% CI 0.778–0.902, P<0.001) and OS (FC: HR 0.806, 95% CI 0.723-0.898, P < 0.001; FCR: HR 0.801, 95% CI 0.709–0.905, P < 0.001) in both FC and FCR treatment groups (Suppl. Table 8). Of note, telomere length included as continuous variable was an independent prognostic factor for OS (HR 0.890, 95% CI 0.805–0.983, P = 0.022) along with FCR (P = 0.009), ECOG  $\ge 1$  (P = 0.020), CIRS score (P = 0.001), 17p- (P = 0.001), unmutated IGHV (P = 0.001)0.043), TP53 mutation (P < 0.001), s-TK > 10 (P = 0.032), and s- $\beta_2$ MG (P = 0.008) (Table 2).





**Fig. 2** Comparison of **a**, **b** PFS (FC: median 27.1 vs. 44.0 months, P < 0.001; FCR: median 44.8 vs. 69.9 months, P < 0.001) and **c**, **d** OS (FC: median 70.9 vs. 89.5 months, P < 0.001; FCR: median 90.2 months vs.

Telomere length associations in the validation cohort CLL4

The telomere length associations identified in the CLL8 were validated using the CLL4 trial cohort. Telomere length in CLL4 was found to be highly variable similar to CLL8, varying from 2.48 kb to 32.41 kb. CLL4 had a higher

median not reached, P = 0.017 between cases with short ( $\leq$  median) and long (>median) telomere length, treated with FC (n = 153 vs. 159) and FCR (n = 157 vs. 151) in CLL8

representation of cases with low and intermediate risk genomic aberration (+12q, normal karyotype and 13q-) compared with CLL8 (73.0% vs. 68.4%, Suppl. Table 1). The longer median telomere length of CLL4 (5.98 kb) may also be owing to differences in DNA extraction method (Qiagen vs Trizol) and DNA quality. Hence, a unifying telomere length cut-point was not used between the two trials for dichotomization and Short telomeres are associated with inferior outcome, genomic complexity, and clonal evolution in...

 Table 2
 Multivariable analysis

 for identification of independent
 prognostic factors in CLL8

 including telomere length as
 continuous variable

| Progression-free survival (PFS) |                         |                               |       | Overall survival (OS) |                                     |                         |                               |       |         |
|---------------------------------|-------------------------|-------------------------------|-------|-----------------------|-------------------------------------|-------------------------|-------------------------------|-------|---------|
| COX<br>regression               | Hazard<br>ratio<br>(HR) | 95%<br>Confidence<br>interval |       | P value               | COX regression                      | Hazard<br>ratio<br>(HR) | 95%<br>Confidence<br>interval |       | P value |
| _                               |                         | Lower                         | Upper |                       |                                     |                         | Lower                         | Upper |         |
| FCR                             | 0.510                   | 0.409                         | 0.635 | < 0.001               | FCR                                 | 0.661                   | 0.485                         | 0.901 | 0.009   |
| TK > 10 (U/L)                   | 1.386                   | 1.062                         | 1.808 | 0.016                 | ECOG≥1                              | 1.465                   | 1.061                         | 2.023 | 0.020   |
| Unmut IGHV                      | 1.724                   | 1.329                         | 2.237 | < 0.001               | CIRS                                | 1.166                   | 1.062                         | 1.280 | 0.001   |
| 17p-                            | 3.441                   | 2.018                         | 5.868 | < 0.001               | Unmut IGHV                          | 1.545                   | 1.014                         | 2.356 | 0.043   |
| 11q-                            | 1.581                   | 1.226                         | 2.038 | < 0.001               | 17p-                                | 2.577                   | 1.476                         | 4.501 | 0.001   |
| TP53 mut                        | 1.881                   | 1.201                         | 2.946 | 0.006                 | TP53 mut                            | 2.640                   | 1.602                         | 4.351 | < 0.001 |
| SF3B1 mut                       | 1.374                   | 1.056                         | 1.787 | 0.018                 | s-TK > 10 (U/L)                     | 1.626                   | 1.042                         | 2.537 | 0.032   |
|                                 |                         |                               |       |                       | s-β <sub>2</sub> MG > 3.5<br>(mg/L) | 1.559                   | 1.121                         | 2.168 | 0.008   |
|                                 |                         |                               |       |                       | Telomere<br>length*                 | 0.890                   | 0.805                         | 0.983 | 0.022   |

\*Continuous variable

analyses. Among the clinical and genetic characteristics, association of short telomeres with high WBC count (P < 0.001), high s-TK (< 0.001) (Table 1), unmutated IGHV (P < 0.001), SF3B1 mutations (P = 0.022), and presence of 11q- (P < 0.001) (Suppl. Table 2) were confirmed in CLL4. Suppl. Fig. 5a, b represent the distribution of telomere length (as continuous variable) in the different gene mutation and genomic aberration subgroups. A full comparison of telomere length associations in CLL4 as grouped and continuous variable is shown in Table 1, Suppl. Table 2 and 7. As in CLL8, all cases with very long telomeres belonged to the favorable prognostic subgroup harboring mutated IGHV and predominantly 13q- as sole abnormality (72%).

Similar to CLL8, short telomeres in CLL4 were significantly associated with shorter PFS in both the treatment arms (F: 23.4 vs. 26.4 months, HR 1.459, 95% CI 1.032-2.062, P = 0.033; FC: 41.0 vs. 51.0 months, HR 1.476, 95% CI 1.012–2.154, P = 0.043; Suppl. Fig. 6a, b). Short telomeres in CLL4 were significantly associated with shorter OS in F but the difference was not statistically significant in FC treatment (F: 77.1 vs. 122.6 months, HR 1.906, 95% CI 1.218–2.983, P = 0.005; FC: 87.6 vs. 110.1 months, HR 1.591, 95% CI 0.998–2.536, P = 0.051; Suppl. Fig. 6c, d). When analyzed as a continuous variable, telomere length showed a significant association with PFS in FC (HR 0.898, 95% CI 0.834–0.968, P = 0.005) and OS in both F (HR 0.819, 95% CI 0.731–0.917, P = 0.001) and FC (HR 0.870, 95% CI 0.786–0.964, P = 0.007) treatment groups (Suppl. Table 8). Multivariable analysis (Suppl. Table 9) including variables that are significant for PFS in univariate analysis (treatment arm, genomic aberrations, mutational status of IGHV and TP53, s-TK and telomere length) showed the FC treatment arm (P = 0.033), 17p- (P < 0.001), 11q- (P < 0.001), TP53 mutation (P = 0.004), and high s-TK (P = 0.003) to be independent prognostic factors for PFS. For OS, when telomere length, age, sex, genomic aberrations by FISH, mutations of IGHV and TP53, s-TK and s- $\beta_2$ MG were included as variables, 17p- (P < 0.001), 11q- (P = 0.003), TP53 mutation (P < 0.001), high s-TK (P = 0.002), unmutated IGHV (P < 0.001), male sex (P = 0.012), and age > 60 (P = 0.041) were significant in multivariable analysis. Telomere length analyzed either as grouped or continuous variable was not an independent prognostic factor for PFS and OS in CLL4.

# Correlation of telomere length with genomic complexity and CLL driver mutations

As telomeres play a key role in maintaining genomic stability, the association of telomere length with genomic complexity was analyzed. The number of copy number alterations (CNA) and loss of heterozygosity (LOH) were identified as a measure of genomic complexity in 353 cases analyzed using SNP array [30]. The sum of CNA and LOH varied from 0 (no detectable alteration) to 9 alterations per case. Genomic complexity was associated with progressive telomere shortening (P < 0.001;  $r_s = -0.411$ ). Also, an increase in genomic complexity was associated with increase in percentage of cases with high-risk genomic aberrations such as 17p-, 11q-, and TP53 mutations (P =0.002;  $R^2 = 0.723$ , Fig. 3a). Similar associations were also identified within the IGHV subgroups (Suppl. Fig. 7a, b). In addition, associations of telomere length and driver mutations were analyzed from WES of 279 patient samples [31]. Similar to the association with genomic complexity, telomere shortening was significantly associated with increase



**Fig. 3 a** Telomere length association with genomic complexity (copy number alterations (CNA) and loss of heterozygosity (LOH)) and high-risk aberrations (17p-, 11q-, and mutated *TP53*). The red lines indicate the median telomere length (left *Y* axis), the green symbols indicate the percentage of high-risk aberrations (right *Y* axis) and the black dots show the distribution of telomere length corresponding to the number of genomic alterations **b**. Telomere length distribution in relation to the number of clonal and subclonal driver events including and **c** excluding genomic aberrations.  $r_s$  in Figs. **a**–**c** is derived from spearman rank correlation and  $R^2$  in **a** from linear regression

in the number of clonal and subclonal driver events, including (P < 0.001;  $r_s = -0.454$ , Fig. 3b) and excluding (P < 0.001;  $r_s = -0.365$ , Fig. 3c) genomic aberrations. Within the IGHV subgroups, the unmutated IGHV cases showed no association between number of driver mutations and telomere length (Suppl. Fig. 7c, d)

# Association of short telomeres with acquisition of poor-risk genetic aberrations and clonal evolution

Among the different genomic aberration subgroups, cases with 17p- and 11q-, associated with *TP53* and *ATM* loss, respectively, had the shortest telomere length (median: 3.4 kb). Therefore, to verify if this presence of short telomeres is a mere consequence of higher proliferation rate of these poor-risk tumors, we compared telomere length with the clone size of these aberrations. Of note, telomere length did not correlate with the clone size of 17p- ( $R^2 < 0.001$ ) and 11q- ( $R^2 = 0.027$ ) as analyzed by FISH. Similarly, telomere length did not correlate with the variant allele frequency of *SF3B1* ( $R^2 = 0.004$ ) and *TP53* mutations (excluding 17p-,  $R^2 = 0.072$ ) analyzed using WES (Fig. 4a–e). Thus, even cases with very low variant allele frequency of these subclonal drivers had very short telomere lengths.

To study if telomere shortening may precede and therefore contribute to clonal evolution in CLL, paired baseline and follow-up samples from 65 cases were analyzed for telomere length dynamics. The median follow-up time for included cases was 44 months (range: 5-83 months). Clonal evolution was present in 24 cases (36.9%) as detected by SNP array analysis. Although there was no significant difference in baseline telomere lengths of cases with and without clonal evolution (P = 0.770), telomere length at follow-up was significantly shortened in cases with clonal evolution (P = 0.003), whereas this was not the case in the group without clonal evolution (P = 0.204; Fig. 4f). The most frequent recurrent genetic alteration acquired upon clonal evolution was 17p- (33.3%) and 41.7% of cases had an acquired 11q- or a 17p- at follow-up. Also, among cases which underwent clonal evolution, 54% had a 17p- or 11qat baseline. In addition, WES analysis of samples at baseline and follow-up identified nine cases that acquired TP53 mutation. Five of nine cases had a pre-existing 17p- or 11qaberration that is known to be associated with short telomeres. Three of four non-high-risk cases with 13q-, 12q +, and normal karyotype had short telomeres (3.28, 3.57, and 4.25 kb, respectively) even before the acquisition of TP53 mutation. The results indicate that in a subset of cases, critical telomere shortening may precede acquisition of high-risk genomic abnormalities.

# Discussion

The relationship between telomere length and cancer biology is highly complex: telomere length of tumors may reflect the telomere of the initially transformed cell, as well as the history of cell divisions since transformation through telomere attrition. Furthermore, telomere shortening may induce cellular adaptation to overcome proliferation **Fig. 4** Telomere length association with percentage of cells with **a** 17p- and **b** 11q- as analyzed using FISH and with mutant allele frequency of **c** *SF3B1* and **d** *TP53* mutations as analyzed using whole-exome sequencing (WES).

e Distribution of telomere length in IGHV, genomic aberration (with their respective cancer cell fractions-CCF, as analyzed by FISH) and gene mutation subgroups as well as association with copy number variation (analyzed by SNP array). f Telomere length at baseline and follow-up in cases with (n = 24) and without clonal evolution (n = 41). Cases with hierarchical 17p- and 11q- are represented as green and brown symbols, respectively. P values and  $R^2$  in **a**–**d** from linear regression and P values in f from Mann-Whitney test



checkpoints. Telomere shortening may accelerate cancer evolution by facilitating acquisition of genomic copy number variation, yielding novel cancer genotypes favoring increased proliferation [11, 12]. Disentangling these complex relationships may be particularly challenging in the context of human cancer, as largely correlative data are available for analysis. Nevertheless, the large size of the studied CLL8 (n = 620) and CLL4 (n = 293) cohorts enables examining several such hypotheses directly in patient data.

First, we considered that telomere length of the cancer cell may correspond to that of the initiating transformed cell. This is of particular importance in CLL, as it tends to affect older individuals [1], and telomere shortening is known to be a part of normal aging [32]. Our data do not support a significant age-related contribution, to the telomere length of the tumor (Table 1). Furthermore, our analysis of age-matched normal B cells, and the CD19subsets show that telomere shortening in CLL does not stem from a defect in the hematopoietic progenitors, but rather seems restricted to the CLL cell fraction (Fig. 1a). Thus, the evolutionary age of the malignant population, reflecting hundreds of divisions after transformation, seems to be more important than pre-transformation aging related shortening. Roth et al. reported significant telomere shortening in T-cells from CLL however the difference in absolute telomere lengths in the T cells from CLL vs. healthy was much smaller in the study. Such a difference could be masked by the limited number of CD19- samples from healthy donors included in the present study. The presence of an association of CD19- telomere length with patient's age indicate that the non-tumor telomere lengths are only minimally affected by the disease [33].

As each cell division results in further telomere attrition, the length of the telomeres may reflect the number of divisions in the cell's past. Indeed, serially sampled cases (median time interval 44 months) demonstrated ongoing telomere shortening (P < 0.001, Fig. 4f) over time. Consistent with this hypothesis, we observed that large tumor burden in the form of higher circulating cell numbers or a more advanced stage is associated with shorter telomeres (Table 1), especially within the more aggressive IGHV unmutated subtype (data not shown). Somatic driver variations associated with adverse outcome (e.g., *SF3B1* mutations) were also linked with shortening, providing further support to the fact that more aggressive disease progression leads to greater telomere attrition (Fig. 1b, c, Suppl. Fig. 5a, b).

On the other hand, when considering the impact of the transformed cell of origin, we found a strong association between longer telomeres and mutated IGHV (Fig. 1b, Suppl. Fig. 5a). This association is well documented [4–9], and can be attributed to the fact that CLL with mutated IGHV are considered to develop from CD5(+), CD27(+), post-germinal center (GC) B-cell subsets [34], which activate telomerase during the GC reaction [35]. Short telomeres in unmutated IGHV may therefore represent an early event in pathogenesis, favoring selection of high-risk clones

in these cases [36], to overcome telomere dysfunction induced genomic instability.

Telomere shortening may also contribute to further diversification through a higher rate of copy number variation, allowing the tumors to more efficiently explore the evolutionary fitness landscape. In support of this notion, we found that shorter telomeres were closely associated with greater number of copy number variations and greater clonal diversity (Fig. 3a-c). In addition, telomere shortening may exert a strong selection pressure for variants such as ATM or TP53 deficiency that will be permissive to ongoing growth despite critical shortening [37]. We have found a close association between these aberrations and shortened telomeres (Fig. 3a). Importantly, this was confirmed even when these mutations were found in small subclones (with relatively small impact on the mean telomere length of the tumor bulk) (Fig. 4a-e), supporting the hypothesis that telomere shortening preceded the appearance of TP53 and ATM defects, and serves as a selection force to favor the emergence of clones bearing these variants. Further studies with multiple follow-up samples from single cases to track emergence of high-risk genomic aberration along with analysis of telomere length dynamics over a longer period of time may be important to confirm the temporal relationship between telomere length and high-risk aberrations.

Collectively, these data sets suggest that telomere erosion in CLL reflects a more-aggressive CLL phenotype, resulting from greater proliferative history, as well as the propensity to increase clonal diversification and specifically select genotypes that will enable further growth despite telomere attrition. Given these factors, it is not surprising that short telomeres in CLL are associated with adverse outcome in both cohorts (Fig. 2, Suppl. Fig. 6), as it may constitute a summation of multiple risk factors such as IGHV unmutated status, advanced stage, chromosomal instability, high-risk genetic alterations, and greater clonal heterogeneity. As telomere shortening seems to be an early event, preceding acquisition of high-risk aberrations, profiling telomere length at diagnosis and through the disease course in nonhigh-risk patients could provide valuable information on disease progression and evolution, thus contributing to the decision of therapy. With the advent of novel non-genotoxic agents such as inhibitors of BTK, PI3K, and BCL2, initial response to therapy is similar across the favorable and unfavorable genetic subgroups [38]. However, identification of complex karyotype as an independent prognostic factor for PFS and OS in CLL treated with ibrutinib [39] may indicate a possible contribution of underlying telomere dysfunction, thus further studies on telomere length associations in the context of novel, non-genotoxic treatments are warranted.

Acknowledgements We thank all patients and their physicians for CLL4 and CLL8 trial participation and donation of samples; Myriam Mendila, Nancy Valente, Stephan Zurfluh, and Jamie Wingate for their support in conception and conduct of the CLL8 trial. This work was supported by the Else Kröner-Fresenius-Stiftung (2010\_Kolleg24), EC (01KT1601, FIRE CLL), BMBF (031L0076C PRECISe), and Deutsche Forschungsgemeinschaft (SFB 1074 projects B1, B2). Genetic analyses in CLL8 were supported by F. Hoffmann-La Roche.

Author contributions B.M.C.J. designed and performed experiments, analyzed and interpreted data and wrote the manuscript; E.T., D.A.L., A.N.T.-W., and J.B. performed experiments, interpreted data and edited the manuscript; J.B. and S.R. analyzed the data and edited the manuscript; S.S. conceived study, designed experiments, interpreted data and wrote the manuscript. The remaining authors collected, analyzed, and interpreted data, and contributed to the manuscript.

#### **Compliance with ethical standards**

**Conflict of interest** M.K.W. and G.F.-R.: employment at F. Hoffmann-La Roche; C.W.: consultancy, membership in board of directors or advisory committees, research funding from Hoffmann-La Roche; K.F.: travel grants from Roche; M.K., U.J., M.H., and S.S.: consultancy, honoraria, and research funding from Roche. The remaining authors declare no conflict of interests.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

### References

- Kipps TJ, Stevenson FK, Wu CJ, Croce CM, Packham G, Wierda WG, et al. Chronic lymphocytic leukaemia. Nat Rev Dis Prim. 2017;3:16096.
- The International CLL-IPI working group. An international prognostic index for patients with chronic lymphocytic leukaemia (CLL-IPI): a meta-analysis of individual patient data. Lancet Oncol. 2016;17:779–90.
- Bechter OE, Eisterer W, Pall G, Hilbe W, Kühr T, Thaler J. Telomere length and telomerase activity predict survival in patients with B cell chronic lymphocytic leukemia. Cancer Res. 1998;58:4918–22.
- Damle RN, Batliwalla FM, Ghiotto F, Valetto A, Albesiano E, Sison C, et al. Telomere length and telomerase activity delineate distinctive replicative features of the B-CLL subgroups defined by immunoglobulin V gene mutations. Blood. 2004;103:375–82.
- Grabowski P, Hultdin M, Karlsson K, Tobin G, Aleskog A, Thunberg U, et al. Telomere length as a prognostic parameter in chronic lymphocytic leukemia with special reference to VH gene mutation status. Blood. 2005;105:4807–12.
- Ricca I, Rocci A, Drandi D, Francese R, Compagno M, Lobetti Bodoni C, et al. Telomere length identifies two different prognostic subgroups among VH-unmutated B-cell chronic lymphocytic leukemia patients. Leukemia. 2007;21:697–705.
- Roos G, Kröber A, Grabowski P, Kienle D, Bühler A, Döhner H, et al. Short telomeres are associated with genetic complexity, high-risk genomic aberrations, and short survival in chronic lymphocytic leukemia. Blood. 2008;111:2246–52.
- Rampazzo E, Bonaldi L, Trentin L, Visco C, Keppel S, Giunco S, et al. Telomere length and telomerase levels delineate subgroups of B-cell chronic lymphocytic leukemia with different biological characteristics and clinical outcomes. Haematologica. 2012;97:56–63.
- 9. Strefford JC, Kadalayil L, Forster J, Rose-Zerilli MJJ, Parker A, Lin TT, et al. Telomere length predicts progression and overall

survival in chronic lymphocytic leukemia: data from the UK LRF CLL4 trial. Leukemia. 2015;29:2411–4.

- Blackburn EH, Greider CW, Szostak JW. Telomeres and telomerase: the path from maize, tetrahymena and yeast to human cancer and aging. Nat Med. 2006;12:1133–8.
- Calado RT, Young NS. Telomere diseases. N Engl J Med. 2009;361:2353–65.
- Xu L, Li S, Stohr BA. The role of telomere biology in cancer. Annu Rev Pathol. 2013;8:49–78.
- Zhang C, Doherty JA, Burgess S, Hung RJ, Lindström S, Kraft P, et al. Genetic determinants of telomere length and risk of common cancers: a Mendelian randomization study. Hum Mol Genet. 2015;24:5356–66.
- Boardman LA, Litzelman K, Seo S, Johnson RA, Vanderboom RJ, Kimmel GW, et al. The association of telomere length with colorectal cancer differs by the age of cancer onset. Clin Transl Gastroenterol. 2014;5:e52.
- Seguí N, Guinó E, Pineda M, Navarro M, Bellido F, Lázaro C, et al. Longer telomeres are associated with cancer risk in MMRproficient hereditary non-polyposis colorectal cancer. PLoS ONE. 2014;9:e86063.
- Baichoo E, Boardman LA. Toward a molecular classification of colorectal cancer: the role of telomere length. Front Oncol. 2014;4:158.
- Wang H, Wang Y, Kota KK, Kallakury B, Mikhail NN, Sayed D, et al. Strong association between long and heterogeneous telomere length in blood lymphocytes and bladder cancer risk in Egyptian. Carcinogenesis. 2015;36:1284–90.
- Du J, Zhu X, Xie C, Dai N, Gu Y, Zhu M, et al. Telomere length, genetic variants and gastric cancer risk in a Chinese population. Carcinogenesis. 2015;36:963–70.
- Svenson U, Roos G. Telomere length as a biological marker in malignancy. Biochim Biophys Acta. 2009;1792:317–23.
- 20. Xu Y, Goldkorn A. Telomere and telomerase therapeutics in cancer. Genes (Basel). 2016;7:pii: E22.
- Fischer K, Bahlo J, Fink AM, Goede V, Herling CD, Cramer P, et al. Long-term remissions after FCR chemoimmunotherapy in previously untreated patients with CLL: updated results of the CLL8 trial. Blood. 2016;127:208–15.
- 22. Eichhorst BF, Busch R, Hopfinger G, Pasold R, Hensel M, Steinbrecher C, et al. Fludarabine plus cyclophosphamide versus fludarabine alone in first-line therapy of younger patients with chronic lymphocytic leukemia. Blood. 2006;107:885–91.
- 23. Hallek M, Fischer K, Fingerle-Rowson G, Fink AM, Busch R, Mayer J, et al. Addition of rituximab to fludarabine and cyclophosphamide in patients with chronic lymphocytic leukaemia: a randomised, open-label, phase 3 trial. Lancet. 2010;376:1164–74.
- 24. Cawthon RM. Telomere measurement by quantitative PCR. Nucleic Acids Res. 2002;30:e47.
- O'Callaghan N, Dhillon V, Thomas P, Fenech M. A quantitative real-time PCR method for absolute telomere length. Biotechniques. 2008;44:807–9.
- Kruk PA, Rampino NJ, Bohr VA. DNA damage and repair in telomeres: relation to aging. Proc Natl Acad Sci USA. 1995;92:258–62.
- Döhner H, Stilgenbauer S, Benner A, Leupolt E, Kröber A, Bullinger L, et al. Genomic aberrations and survival in chronic lymphocytic leukemia. N Engl J Med. 2000;343:1910–6.
- Kröber A, Seiler T, Benner A, Bullinger L, Brückle E, Lichter P, et al. V(H) mutation status, CD38 expression level, genomic aberrations, and survival in chronic lymphocytic leukemia. Blood. 2002;100:1410–6.
- 29. Stilgenbauer S, Schnaiter A, Paschka P, Zenz T, Rossi M, Döhner K, et al. Gene mutations and treatment outcome in chronic lymphocytic leukemia: results from the CLL8 trial. Blood. 2014;123:3247–54.

- Edelmann J, Holzmann K, Miller F, Winkler D, Bühler A, Zenz T, et al. High-resolution genomic profiling of chronic lymphocytic leukemia reveals new recurrent genomic alterations. Blood. 2012;120:4783–94.
- Landau DA, Tausch E, Taylor-Weiner AN, Stewart C, Reiter JG, Bahlo J, et al. Mutations driving CLL and their evolution in progression and relapse. Nature. 2015;526:525–30.
- Spivak IM, Mikhelson VM, Spivak DL. Telomere length, telomerase activity, stress, and aging. Adv Gerontol. 2016;6:29–35.
- Röth A, de Beer D, Nückel H, Sellmann L, Dührsen U, Dürig J, et al. Significantly shorter telomeres in T-cells of patients with ZAP-70<sup>+</sup> /CD38<sup>+</sup> chronic lymphocytic leukaemia. Br J Haematol. 2008;143:383–6.
- Seifert M, Sellmann L, Bloehdorn J, Wein F, Stilgenbauer S, Dürig J, et al. Cellular origin and pathophysiology of chronic lymphocytic leukemia. J Exp Med. 2012;209:2183–98.
- Weng NP, Granger L, Hodes RJ. Telomere lengthening and telomerase activation during human B cell differentiation. Proc Natl Acad Sci USA. 1997;94:10827–32.

- 36. Jeromin S, Weissmann S, Dicker F, Kern W, Schnittger S, Kohlmann A, et al. IGHV mutation status does not add prognostic information in the background of mutations in TP53 and SF3B1, and deletions of 17p and 11q, which are independent adverse prognostic parameters In CLL. Blood. 2013; 122:4110.
- 37. Herbig U, Jobling WA, Chen BPC, Chen DJ, Sedivy JM. Telomere shortening triggers senescence of human cells through a pathway involving ATM, p53, and p21(CIP1), but not p16 (INK4a). Mol Cell. 2004;14:501–13.
- Stilgenbauer S. Prognostic markers and standard management of chronic lymphocytic leukemia. Hematol Am Soc Hematol Educ Program. 2015;2015:368–77.
- 39. Thompson PA, O'Brien SM, Wierda WG, Ferrajoli A, Stingo F, Smith SC, et al. Complex karyotype is a stronger predictor than del(17p) for an inferior outcome in relapsed or refractory chronic lymphocytic leukemia patients treated with ibrutinib-based regimens. Cancer. 2015;121: 3612–21.